


JANUARY 2022, Volume 10, Issue 01



ICAR - National Institute of Veterinary
Epidemiology and Disease Informatics
(ICAR - NIVEDI)



2022
March

LIVESTOCK DISEASE FOREWARNING BULLETIN

Powered by Artificial Intelligence

PUBLISHED BY:
DIRECTOR
ICAR-NIVEDI

©2022 ICAR-NIVEDI

NOW LINKED WITH
FARMER REGISTRATION AND UNIFIED
BENEFICIARY INFORMATION SYSTEM
(FRUITS)



Citation: Suresh K P, Hemadri D, Patil S S, Krishnamoorthy P, Siju S J and Shome B R. Livestock Disease Risk Forewarning Bulletin - March 2022, ICAR-NIVEDI, Bengaluru, 10(01): 1-126.

Month & Year: January, 2022

Published by: Director, ICAR- National Institute of Veterinary Epidemiology and Disease Informatics (NIVEDI), Yelahanka, Bengaluru-560064.

PME NUMBER: NIVEDI/PMEC/Forewarning Bulletin/2021-22/06

©ICAR-NIVEDI

Prepared By: Dr. K. P. Suresh
Dr. D. Hemadri
Dr. S.S. Patil
Dr. P. Krishnamoorthy
Dr. S.J. Siju

Printed by
Cnu Graphics Printers
35/1, S End Rd, Jai Bheema Nagar, Malleswara, Bengaluru,
Karnataka 560003

Front Page Design by: Ms. Apoorva Hemadri

Disclaimer

The forewarnings are based on the retrospective disease data available in the NADRES database. Hence, for those states wherein data is limited/less, the forewarning may not be realistic. Further the forewarning will not take into consideration the control measures that are *in situ*.

Acknowledgement

I would like to acknowledge the constant support and inspiration from Hon'ble Secretary, DARE and Director General, ICAR, Government of India, New Delhi.

I would like to express sincere everlasting gratitude to Hon'ble Deputy Director General (Animal Science) for his constant encouragement, support and guidance.

I would also like to express sincere gratitude to Department of Animal Husbandry and Dairying (DAHD), Ministry of Agriculture and Farmers Welfare, Government of India for providing the livestock population data for preparation of this bulletin.

Animal Husbandry Departments of state governments and also AICRP on ADMAS centres are gratefully acknowledged for the timely submission of reports of livestock disease outbreak data. I am thankful to all the scientific and technical staff of ICAR-NIVEDI for their feedback and support. I sincerely acknowledge the Statistical Division of DAHD for providing the data on livestock census.

Furthermore, I would also like to acknowledge with much appreciation the crucial role of Dr. K. P. Suresh, Principal Scientist and support received from the scientists, Dr. D. Hemadri, Dr. S.S. Patil, Dr. P. Krishnamoorthy, Dr. S.J. Siju and SRF's, JRF's, Young Professionals and other contractual staff working in Disease Informatics Lab/Spatial Epidemiology Lab in preparing this monthly bulletin.

Director (Acting)
ICAR- NIVEDI

TABLE OF CONTENT

| Sl No | Content | Page Number |
|--------------|---|--------------------|
| 1 | About the Bulletin | 1 |
| 2 | Summary of the Forewarning Bulletin | 2 |
| 3 | Introduction to NADRES v2 | 4 |
| 4 | Forewarning Methodology | 6 |
| | I. Material | 6 |
| | II. NADRES v2 Data flow and AI Based Data capturing diagram | 8 |
| | II .Weighted outbreak score | 9 |
| | IV. Forecasting of weather parameters | 9 |
| | V. Implementation of Principal Component Analysis | 11 |
| | VI. Machine Learning Models | 11 |
| 5 | Accuracy of Prediction | 12 |
| 6 | Moran's I for clustering of Livestock diseases | 13 |
| 7 | R Software | 13 |
| 8 | Forewarning of livestock disease for the month of March, 2022 | 14 |
| | I. District wise Livestock Disease Risk Forewarning | 14 |
| | II. Glimpse of Predicted disease outbreaks | 60 |
| | III. Diseases, Species affected, clinical signs and its preventive measures | 73 |
| | IV. Risk Prediction -Livestock disease forewarning maps | 78 |
| | V. Forecasting of remote sensing and meteorological parameters | 91 |
| | VI. Significant Weather Parameters Table | 93 |
| 9 | Post prediction Validation | 94 |
| | I. Correlational Assessment | 100 |
| 10 | Launch of Mobile Android app. & link to download | 102 |
| 11 | Farmers Empowerment through IT: Disease Risk Communication | 104 |
| 12 | Appendix | 105 |
| | a). R Code | 105 |
| | b). Abbreviations | 111 |
| 13 | Infection with SARS-COV-2 in animals | 112 |
| | a). Epidemiology of COVID-19 | 118 |
| 14 | Customer/Client Feedback Form | 125 |

1. About the Bulletin...

Livestock sector plays a crucial role in the rural economy of India as around 20.5 million people depend upon livestock for their livelihood. Even though the investment in the livestock sector is meagre, tremendous achievements have been observed in the sector during the last decade. As it is an important component in poverty alleviation programmes, continuous emphasis is being laid on this sector for enhancing the quality of the primary and secondary products in the international market, which in turn demands improved animal health. Therefore, livestock development programmes cannot succeed unless a well-organized animal health service is built up and in place for safeguarding the livestock against economically important diseases.

India has made a noteworthy success in the eradication of Rinderpest (RP), CBPP, AHS and Dourine. However, there are several other infectious and non-infectious diseases prevailing in the country causing huge annual economic loss. Prevention, control and eradication of the animal diseases need a thorough understanding of the epidemiology as well their economic impact.

National Institute of Veterinary Epidemiology and Disease Informatics (NIVEDI) has the mandate to carry out research activities in the area of veterinary epidemiology and disease informatics. With the eradication of RP successfully, India has not only proved its ability to face the challenges but also to succeed, despite various limitations. Similar efforts are needed to control and eradicate diseases like FMD, PPR, Brucellosis, CSF, HS etc., which cause huge economic loss annually to the livestock industry. To this end, ICAR-NIVEDI has identified 13 priority diseases, based on the past incidence patterns and has built a strong database of these diseases. The database, which is the backbone of the National Animal Disease Referral Expert System (NADRESv2), is used for providing monthly livestock disease forewarning, which is compiled in this monthly bulletin to alert the animal husbandry departments, both at the National/state level, to take appropriate control measures. We hope users/stakeholders find this bulletin useful in their quest to control livestock diseases. This forewarning bulletin will assist the field Veterinarians in adopting appropriate preventive and control measures, thereby reducing the occurrence of livestock disease outbreaks.

2. SUMMARY OF THE FOREWARNING BULLETIN....

The association between infectious diseases and the climate was known from ancient times. Hippocrates observed in the 5th century that epidemics were associated with natural phenomenon rather than divinities or demons. In modern times, our increasing capabilities to detect and predict climate variations joined with growing evidence for global climate change, have powered interest in understanding the impacts of climate on animal health, particularly the emergence and transmission of infectious disease agents. Simple reasoning suggests that climate can affect infectious disease patterns because disease agents (viruses, bacteria, and parasites) and their vectors are sensitive to temperature, moisture, and other ambient environmental conditions.

India being an agriculture-based country, the livestock sector plays a vital role in contributing to the economy. A robust reporting and forewarning system enable the concerned authorities in disease preparedness and awareness of the risk associated with livestock disease. Therefore, the economic loss due to morbidity and mortality of the animals is reduced thereby helps to increase the productivity in terms of egg, meat, and dairy products. National Animal Disease Referral Expert System database is a weather-based forewarning system enabled with an artificial intelligence system developed by ICAR- National Institute of Veterinary Epidemiology & Disease Informatics Bengaluru, Karnataka state, India that forecast potential threats from pathogens two months in advance to provide the stakeholders with sufficient timeline for awareness and preparedness to act. Artificial Intelligence (AI) and Machine Learning (ML) models use the programmed algorithms that receive and analyse input data to predict output (Infectious risk prediction) values within an acceptable range. As new data fed into these algorithms, they learn and optimize their operations to improve performance, developing intelligence over time.

SUMMARY

The livestock disease forecasting for March 2022 revealed Jharkhand (131), Uttar Pradesh (77), Karnataka (51), Assam (46) and West Bengal (39) as the top five states with high predicted livestock disease outbreaks.

Among the predicted diseases, control programmes are in full swing for FMD and PPR in the country and due attention is demanded by the predicted disease outbreaks of these diseases. Among the expected disease outbreaks, the predicted FMD outbreaks are more in Kerala (13) followed by Karnataka (12) and Jharkhand (10) whereas predicted PPR outbreaks are more in Jharkhand (9) and West Bengal (9). Further, the co-occurrence of FMD and HS can be expected in Haryana, Jharkhand, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Mizoram, Odisha, Rajasthan, Tamil Nadu, Tripura and West Bengal. Among the different diseases in livestock, the predicted outbreaks are expected to be high for FMD (63) and Babesiosis (61).

The major challenges for the effective disease control programme being the lack of thorough understanding about the complexity of disease dynamics, wide host range of pathogens, widening of niche of pathogens due to climate change etc. Further, India has witnessed emergence and re-emergence of various infectious pathogens during the last decade, of which most of the diseases are of zoonotic in nature which urge for the necessity of strengthening of monitoring and surveillance system in the country. Effective control programme for major livestock diseases in the country can be efficiently addressed by planning and execution of available control measures in the high-risk areas and routine surveillance and monitoring of diseases.

Table S1. Summary of State wise Livestock Disease forewarning for March, 2022

| Sl. No | State Name | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis | Total number of diseases predicted |
|---|------------------------|-----------|------------|-----------|----------|-----------|-------------|-----------|-----------|-----------|-----------|-----------|--------------|----------------|------------------------------------|
| 1 | Andaman and Nicobar | 0 | 0 | 0 | 0 | 0 | 3 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 04 |
| 2 | Andhra Pradesh | 3 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 1 | 1 | 0 | 0 | 0 | 07 |
| 3 | Arunachal Pradesh | 0 | 0 | 0 | 0 | 0 | 4 | 1 | 0 | 0 | 3 | 2 | 0 | 0 | 10 |
| 4 | Assam | 0 | 2 | 12 | 0 | 2 | 12 | 0 | 5 | 6 | 1 | 5 | 1 | 0 | 46 |
| 5 | Bihar | 0 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 2 | 06 |
| 6 | Chandigarh | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 00 |
| 7 | Chhattisgarh | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 01 |
| 8 | Dadra and Nagar Haveli | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 00 |
| 9 | Daman and Diu | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 00 |
| 10 | Goa | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 02 |
| 11 | Gujarat | 0 | 0 | 0 | 0 | 1 | 0 | 2 | 0 | 2 | 1 | 0 | 0 | 0 | 06 |
| 12 | Haryana | 0 | 0 | 0 | 0 | 2 | 0 | 1 | 2 | 3 | 0 | 1 | 1 | 1 | 11 |
| 13 | Himachal Pradesh | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 1 | 1 | 0 | 0 | 0 | 04 |
| 14 | Jammu and Kashmir | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 10 | 0 | 0 | 0 | 10 |
| 15 | Jharkhand | 0 | 23 | 2 | 0 | 8 | 22 | 10 | 4 | 9 | 2 | 5 | 23 | 23 | 131 |
| 16 | Karnataka | 7 | 0 | 6 | 0 | 5 | 0 | 12 | 3 | 4 | 14 | 0 | 0 | 0 | 51 |
| 17 | Kerala | 3 | 2 | 0 | 0 | 0 | 0 | 13 | 2 | 4 | 0 | 2 | 3 | 0 | 29 |
| 18 | Lakshadweep | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 00 |
| 19 | Madhya Pradesh | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 03 |
| 20 | Maharashtra | 0 | 0 | 1 | 0 | 1 | 0 | 3 | 2 | 6 | 3 | 0 | 3 | 0 | 19 |
| 21 | Manipur | 0 | 0 | 4 | 0 | 1 | 5 | 0 | 0 | 0 | 2 | 4 | 0 | 0 | 16 |
| 22 | Meghalaya | 0 | 0 | 6 | 0 | 0 | 0 | 4 | 3 | 0 | 0 | 5 | 0 | 0 | 18 |
| 23 | Mizoram | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 2 | 7 | 0 | 0 | 11 |
| 24 | Nagaland | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 01 |
| 25 | NCT of Delhi | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 01 |
| 26 | Odisha | 1 | 1 | 3 | 0 | 0 | 0 | 2 | 1 | 5 | 0 | 0 | 1 | 0 | 14 |
| 27 | Puducherry | 0 | 2 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 04 |
| 28 | Punjab | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 01 |
| 29 | Rajasthan | 0 | 2 | 0 | 0 | 2 | 1 | 2 | 4 | 4 | 0 | 1 | 0 | 0 | 16 |
| 30 | Sikkim | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 04 |
| 31 | Tamil Nadu | 3 | 0 | 2 | 0 | 0 | 0 | 1 | 1 | 3 | 3 | 1 | 0 | 0 | 14 |
| 32 | Telangana | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 02 |
| 33 | Tripura | 1 | 0 | 3 | 0 | 0 | 3 | 2 | 3 | 0 | 3 | 3 | 0 | 0 | 18 |
| 34 | Uttar Pradesh | 0 | 21 | 0 | 0 | 1 | 5 | 2 | 0 | 1 | 0 | 0 | 18 | 29 | 77 |
| 35 | Uttarakhand | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 01 |
| 36 | West Bengal | 0 | 3 | 7 | 0 | 0 | 1 | 3 | 4 | 9 | 4 | 0 | 7 | 1 | 39 |
| Total number of districts likely to report | | 19 | 61 | 48 | 0 | 24 | 57 | 63 | 38 | 60 | 52 | 40 | 59 | 56 | 577 |

*Number of predicted disease incidence was summarised considering only High risk and Very high risk

3. INTRODUCTION TO NADRES v2

The geographic and seasonal distribution of many infectious diseases are associated with climate and therefore the possibility of using seasonal climate forecasts as predictive indicators in disease early warning system (EWS) became imminent. In this context, ICAR-NIVEDI, in its quest for achieving better livestock health, had developed an interactive web portal named "National Animal Disease Referral Expert System (NADRES)" during early part of the first decade of the millennium. The web portal, which was developed from the financial support of National Agricultural Technology Project, was launched in the year 2005. The portal which is interactive, allows the user/stakeholder to access livestock disease forewarning ($n=13$) at the district level for entire country two months in advance. The portal which was initially built on oracle platform was later changed to MySQL platform to store the administrator provided disease information and other relevant meteorological and risk factor information. However, with the availability of remote sensed satellite images and the advancement in information technology and statistical algorithms, the upgradation of NADRES became inevitable. To this end, a newer version of NADRES (NADRES V2) has been developed and is ready for release.

How it is different from previous version?

In brief, it can be said that NADRES v2 underwent a sea change not only in its internal structure but also in its physical design. As a result, now the central menu bar consists of Home, About us, Risk factors, Analysis, Livestock disease, post prediction validation and contact details. Risk factors menu comprises of details on resolution, time interval, units and source of 11 meteorological and 5 remote sensing parameters. Analytics menu has various analysis options. The newly created livestock disease menu has the details regarding species affected, clinical signs and preventive measures to be adopted for the 13 economically important diseases. Post prediction validation menu contains the outbreak reports vs prediction. The menu bar on the RHS tabs include online GIS, state wise Livestock disease forecast, district wise Livestock disease forecast, Epi-calculator, download links for mobile app, etc. The website now hosts disease maps in the form of choropleth maps for 13 diseases in two time periods (1990-2000 and 2000-2018). Similarly, disease trends plots exhibit periodic regression plots providing future trend for the disease. On the LHS, Login menu is provided for authorized persons to login and enter disease details and other related parameters. Disease maps provide choropleth maps for 13 diseases in two time periods (1990-2000 and 2000-2018) is presented. Disease trends- Periodic regression plots are exhibited for prediction of the diseases. Auto-messaging option has been created to send the reminders in the form of text messages to concerned PI's and Co-PI's of AICRP centers for submission of outbreak reports. This message is sent weekly to all the concerned officials. Additionally, a message is sent to the concerned veterinary officers in Karnataka for initiation of preventive measures for the forewarned diseases at the block level. Plans are in place to incorporate farmers' and local vets' mobile numbers in to the list so that they may be asked to initiate preventive measures for the forewarned diseases.



Fig 3.1. NADRES V₂ Home page

The forewarning methodology used is unique and has not been used earlier for livestock disease forewarning in India. Following few paragraphs describe about the forewarning methodology used. It is a well-known fact that weather plays an important role in the precipitation of many diseases and therefore, the climatic parameters such as land surface temperature (LST), precipitation, wind velocity, humidity etc are considered as risk parameters. These parameters along with other non-climatic parameters such as livestock population, density, Normalized Differential Vegetation Index (NDVI), soil moisture constitute the overall risk parameters. A total of 24 such parameters are collected/generated at village level and then aggregated to district level before these are used for analysis.

In addition to the output provided at interactive web portal, the NADRES output are also published in the form of monthly livestock disease forewarning bulletins. The prediction results come with a disclaimer that forewarnings do not take into account of the control measures that already in situ and also may not be realistic for those regions where the data is either unavailable or limited. This bulletin provides the likely occurrence of the 13 shortlisted diseases two months in advance at the district level, disease forewarning maps, prediction accuracy, details on diseases, species affected, clinical signs and its preventive measures.

In summary, it can be said that NADRES v₂ has underwent substantial changes not only in its internal structure but also in its physical design and can be a useful tool for visitors of the website, farmers, vets, policy makers etc.

4. Forewarning Methodology

I.Materials

Livestock disease data

Previous 10 years' livestock disease outbreak data retrieved from the NADRES database linked with Risk factors data.

Livestock population data

The population data at village level for five major livestock species viz., cattle, buffalo, sheep, goat and pigs were obtained from 20th Livestock census (2019) from Department of statistics, DAHD, GOI.

| Species-wise & Category-wise Livestock Population (in thousands) | | | | | |
|---|-----------------|------------|--------------------|--------------------|------------|
| Sl No | Species | Category | Population in 2012 | Population in 2019 | % Change |
| 1 | Cattle | Exotic | 39732 | 51356 | 29.3 |
| | | Indigenous | 151172 | 142106 | -6 |
| | | Total | 190904 | 193462 | 1.3 |
| 2 | Buffalo | Total | 108702 | 109852 | 1.1 |
| 3 | Sheep | Exotic | 3781 | 4088 | 8.1 |
| | | Indigenous | 61288 | 70172 | 14.5 |
| | | Total | 65069 | 74260 | 14.1 |
| 4 | Goat | Total | 135173 | 148885 | 10.1 |
| 5 | Pig | Exotic | 2456 | 1897 | -22.8 |
| | | Indigenous | 7837 | 7159 | -8.7 |
| | | Total | 10293 | 9056 | -12 |
| 6 | Yaks | Total | 77 | 58 | -24.7 |
| 7 | Mithuns | Total | 298 | 386 | 29.5 |
| 8 | Horses & Ponies | Total | 625 | 342 | -45.3 |
| 9 | Mules | Total | 196 | 84 | -57.1 |
| 10 | Donkeys | Total | 319 | 124 | -61.1 |
| 11 | Camels | Total | 400 | 252 | -37 |
| Total Livestock | | | 512056 | 536761 | 4.8 |

Meteorological and Remotely Sensed Data:

The parameters such as air temperature ($^{\circ}\text{C}$), perceptible water (mm), pressure (millibar), relative humidity (%) and sea level pressure (millibar) were extracted from National Centre for environmental prediction (NCEP). The parameters such as potential evapotranspiration (PET), Enhanced Vegetation Index (EVI), Leaf Area Index (LAI), Land Surface Temperature (LST), Normalised Difference Vegetation Index (NDVI) were extracted from remote sensed images from MODIS website (<https://modis.gsfc.nasa.gov/>). In brief, the MODIS products from NASA-TERRA satellite was downloaded for the Indian locations by specifying the tiles (H24V5, H25V6, H24V6, H24V7, H25V7, H25V8, H26V7, H26V6) from 2001 to till date.

The details are given below;

| PRODUCT | Science Data Sets (HDF Layers) |
|-----------------|--|
| MOD15A2H | Lai_500m(Leaf area index) 8 days average |
| MOD16A2 | PET_500m (Total Potential Evapotranspiration) 8 days average |
| MOD11A2 | LST_Day_1km (Daytime Land Surface Temperature) 8 days average |
| MOD13A1 | <ul style="list-style-type: none"> i. 500m 16 days NDVI (Normalized Difference Vegetation Index) ii. Enhanced Vegetation Index (EVI) 16 days average |

The downloaded HDF files (Datasets, which are multidimensional arrays (layers) of a homogeneous type) were converted to GeoTIFF files (single layer data) using R packages, which were later used to extract the parameters by linking it with the sinusoidal values of the Indian villages. The scale factors were multiplied for the extracted values as specified by the MODIS data products to get the values of the parameters. As shown above, the atmospherically corrected NDVI was collected on 16-day interval at 250-meter resolution using MODIS product MOD13A1 and LST was collected on 8-day interval using MOD11A2 at 1 KM resolution.

The parameters such as rainfall, soil moisture and wind speed were obtained from Global Land Data Assimilation System of NASA (<https://disc.gsfc.nasa.gov>). The remaining parameters were downloaded from climatic research unit (CRU) of University of East Anglia website. It is worth mentioning that the entire process of extraction, assimilation, processing and aligning have been done using R programming language and R environment. After aligning the climatic and non-climatic data with the disease and the livestock population data (aggregated at the district level), the statistical analysis was performed in the R environment.

Initially, two regression models and six machine learning models were applied to test their suitability to fit the data and in all, three models; one regression model (**Generalized Linear Model (GLM)**) and two machine learning models, *viz.*, Gradient Boosting Machine Learning Algorithm (**GBM**) and Random Forest (**RF**), which fitted to data well were incorporated for the purpose of disease prediction. The models were trained using the case and control data available at ICAR-NIVEDI. Validation of the models were done by dividing the total observations for a particular disease into marker samples and validation samples and accuracy was tested in terms of discrimination power, which was done using Receiving Operating Characteristics (**ROC**), Cohen Kappa (**Heildke Skill Score**) and True Skill statistics (**TSS**). Once the models produce the probability value, it was used for categorizing the risk. Briefly, when all the models produce the p value of more than 0.5, then the highest p value is used for determining the high-risk category. If all the models or any one model produces the p value of less than 0.5, then the lowest p value was used for categorizing lower risk. This was done to minimize the false alert. Thus, the risk predictions based on the probability values ranging from 0-1 are made as follows; Very High Risk ($p=0.81-1.0$), High Risk ($p=0.61-0.80$), Moderate Risk ($p=0.41-0.60$), Low Risk ($p=0.21-0.40$), Very Low Risk ($p=0.0-0.20$) and No Risk ($p=0.0$) for the occurrence of a said disease. It is believed that categorizing districts in to various risk categories will help the stake holders to effectively utilize the available resources (money and manpower).

II.NADRES v2 Data Flow and Data Processing Diagram

A) Data Flow Diagram:





Fig 4.1. NADRES v2 Data Flow Diagram.

B) Artificial Intelligence enabled Data Capturing and Forewarning System:



III. Weighted Outbreak Score

The outbreak data for the month of forecasting is extracted from NADRES database for the period of 10 years from current year. Outbreak data of 13 important livestock diseases are considered. The data is aggregated at district level and the weighted score is defined based on the number of outbreaks for each district in each month considering last 10 years. The weightage score was assigned as 0 for less than three number of outbreaks in the last 10 years for selected month, score 1 for 3–6 number of outbreaks and 2 for more than 6 outbreaks. This weightage score for each district is labelled as risk variable in building the models and risk maps.




Fig 4.3. Top ten livestock diseases (2020)

IV. Forecasting of Weather Parameters

Weather forecasting has been one of the most challenging problems around the world because of both its practical value in meteorology and popular sphere for scientific research. Weather forecast systems are among the most complex equation systems that computer has to solve. A great quantity of data, coming from satellites, ground stations and sensors located around our planet send daily information that must be used to foresee the weather situation in next hours and days all around. Weather forecasts provide critical information about future weather. There are various techniques involved in weather forecasting, from relatively simple observation of the sky to highly complex computerized mathematical models. Further, forecast products by Indian Metrological department were used for validation of our forecasts (https://mausam.imd.gov.in/imd_latest/contents/extendedrangeforecast.php).

Following are the basic steps of forecasting process:

1. Determine the forecast's purpose
2. Establish a time horizon
3. Select a forecasting technique
4. Gather and analyse data
5. Perform the forecast
6. Monitor the forecast and use it in prediction of disease

Statistical Models used for forecasting of weather and remotely sensed variables

ARIMA stands for Autoregressive Integrated Moving Average. ARIMA is also known as Box-Jenkins approach. Box and Jenkins claimed that non-stationary data can be made stationary by differencing the series, Y_t . The general model for Y_t is written as,

$$Y_t = \phi_1 Y_{t-1} + \phi_2 Y_{t-2} \dots \phi_p Y_{t-p} + \epsilon_t + \theta_1 \epsilon_{t-1} + \theta_2 \epsilon_{t-2} + \dots \theta_q \epsilon_{t-q}$$

Where, Y_t is the differenced time series value, ϕ and θ are unknown parameters and ϵ are independent identically distributed error terms with zero mean. Here, Y_t is expressed in terms of its past values and the current and past values of error terms.

The ARIMA Model combines three basic Methods:

- Auto Regression (AR) – In auto-regression the values of a given time series data are regressed on their own lagged values, which is indicated by the "p" value in the model.
- Differencing (I-for Integrated) – This involves differencing the time series data to remove the trend and convert a non-stationary time series to a stationary one. This is indicated by the "d" value in the model. If $d = 1$, it looks at the difference between two-time series entries, if $d = 2$ it looks at the differences of the differences obtained at $d = 1$, and so forth.
- Moving Average (MA) – The moving average nature of the model is represented by the "q" value which is the number of lagged values of the error term.

This model is called Autoregressive Integrated Moving Average or ARIMA (p, d,q) of Y_t . We will follow the steps enumerated below to build our model. ARIMA models were run in 18 combinations of p, d, q. Based on the minimum AIC value, the order of ARIMA model was selected. This order was used for the prediction of all the weather parameters used in developing disease forewarning models.

V. Implementation of Principal Component Analysis

Large datasets are gradually common and are often difficult to interpret. Principal Component Analysis (PCA) is a technique for reducing the dimensionality of such datasets, increasing the interpretability but at the same time, minimizing the information loss. The PCA is employed in NADRES v2 by creating new uncorrelated variables that successively maximize the variance. This means that ` preserving as much variability as possible` translates into finding new variables that are linear functions of those in the original dataset, that successively maximize variance and that are uncorrelated with each other. Determining such new variables, the principal components (PCs) reduce to solve an eigenvalue/eigenvector problem. PCA can be based on either covariance matrix or the correlation matrix and the main use of PCA are descriptive.

In the present study, all the meteorological and remote sensing variables are considering for PCA, with correlation matrix, the final output of principal components which are independent of each were considered for further ML modelling and risk estimation.

VI. Machine Learning Models

Disease outbreak data were aligned with generated risk variables to the respective latitude and longitude, which were subjected to climate-disease modelling. A number of models were fit to aligned data and tested for accuracy in terms of discrimination power. Two regression models, Generalized Linear Models (GLM) and Generalized Additive Models (GAM) and six machine learning algorithms, i.e. Random Forest (RF), Boosted Regression Tree (BRT), Artificial Neural Network (ANN), Multiple Adaptive Regression Spline (MARS), Flexible Discriminant Analysis (FDA) and Classification Tree Analysis (CTA) were employed for disease modelling. Different modelling methods return different types of 'model object' and all these model objects could be used for the predict function to make predictions for any combinations of values of independent variables. Response plots were created to explore and understand model predictions.

The fitted models were assessed for their discriminating power using Receiving Operating Characteristic (ROC) curve, Cohen's Kappa (Heildke Skill Score) and True Skill Statistics (TSS). These measures were used to evaluate the quality of predictions based on presence-absence data. Raster Stack was used to combine the results of individual predictions by different model methods. All the models were assessed for overfitting.

The outcome of best fitted models was in probability of disease occurrence and was categorised into 6 risk levels as No risk (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR) and Very high risk (VHR) for enabling the stakeholders to take appropriate control measures by suitably allocating available resources.

5. Accuracy of Prediction

| Serial No. | Diseases | Accuracy (%) |
|------------|----------------------------|--------------|
| 1. | Anthrax | 99.69 |
| 2. | Babesiosis | 97.69 |
| 3. | Black Quarter | 99.23 |
| 4. | Blue Tongue | 99.07 |
| 5. | Enterotoxaemia | 99.85 |
| 6. | Fasciolosis | 99.69 |
| 7. | Foot and mouth disease | 94.14 |
| 8. | Haemorrhagic septicaemia | 98.15 |
| 9. | Peste des Petits Ruminants | 96.60 |
| 10. | Sheep & Goat pox | 97.99 |
| 11. | Swine fever | 98.30 |
| 12. | Theileriosis | 98.61 |
| 13. | Trypanosomosis | 97.53 |

Aggregation and prediction of livestock diseases at district level leading to higher accuracy.

- **Formula Used:** The **Accuracy** of disease prediction was calculated using the following formula.

$$\frac{TP + TN}{Total} * 100$$

TP-True Positive Observations, TN-True Negative Observations, Total- Total observations.

- Internal Accuracy was performed using 10 years of data. Accuracy obtained was >90% for all the diseases predicted.
- Despite the power of climate and disease risk models, considerable uncertainties remain, identifying these uncertainties, highlighting importance of improved data may improve the model accuracy, realism, confidence, together with translating uncertainties in model inputs into uncertainties in model outputs, are important benefits of modelling.

6. Moran's I for clustering of Livestock diseases

Moran's I is a tool that measures spatial autocorrelation (feature similarity) based on both feature locations and feature values simultaneously. Given a set of

features and an associated attribute, it evaluates whether the pattern expressed is clustered, dispersed, or random. The tool calculates the Moran's I Index value and both a Z score and p-value evaluating the significance of that index. In general, a Moran's Index value near +1.0 indicates clustering while an index value near -1.0 indicates dispersion.

Autocorrelation tool, the null hypothesis states that "there is no spatial clustering of the values associated with the geographic features in the study area ." When the p-value is small and the absolute value of the Z score is large enough that it falls outside of the desired confidence level, the null hypothesis can be rejected . If the index value is greater than 0, the set of features exhibits a clustered pattern .If the value is less than 0, the set of features exhibits a dispersed pattern.

7. R Software

R is a programming language and software environment for statistical analysis, graphics representation and reporting. R is a simple and effective programming language which includes conditionals, loops, user defined recursive functions and input and output facilities. R statistical software version 3.1.3 (version 3.4.3, R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>) was used as an integrated suite for data mining, calculation and graphical display. Several R packages like *openxlsx*, *raster*, *RMySQL*, *rgdal*, *RColorBrewer*, *sqldf*, *sp*, *spdep*, *xlsx*, *plyr*, *randomFores*, *dismo*, *SDMTool*, *dplyr*, *tmap* and *data.table* were used for data extraction, data alignment, annotation, analysis, modelling and risk mapping.

8. Forewarning of Livestock Disease for the Month of March, 2022

i). District wise Livestock Disease forewarning:

District wise Livestock Disease Risk Forewarning for March, 2022: Andaman and Nicobar

| Districts of Andaman and Nicobar | Livestock Diseases | | | | | | | | | | | | |
|----------------------------------|--------------------|------------|----|----|----|-------------|-----|----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Nicobars | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR | NR |
| North & Middle Andaman | NR | NR | NR | NR | NR | VHR | VLR | NR | NR | NR | VLR | NR | NR |
| South Andaman | NR | NR | NR | NR | NR | VHR | VLR | MR | NR | HR | VLR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Andhra Pradesh

| Districts of Andhra Pradesh | Livestock Diseases | | | | | | | | | | | | |
|-----------------------------|--------------------|------------|------------|-----|----|-------------|-----|------------|------------|------------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Anantapur | VHR | NR | NR | LR | NR | NR | NR | VHR | NR | NR | NR | NR | NR |
| Chittoor | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| East Godavari | NR | NR | VHR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Guntur | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR | NR | NR | NR |
| Krishna | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Kurnool | VHR | NR | NR | LR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Prakasam | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Sri Potti Sriramulu Nellore | NR | NR | NR | VLR | NR | NR | VLR | NR | VHR | VHR | NR | NR | NR |
| Srikakulam | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Visakhapatnam | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Vizianagaram | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| West Godavari | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Y.S.R. | VHR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Arunachal Pradesh

| Districts of Arunachal Pradesh | Livestock Diseases | | | | | | | | | | | | |
|--------------------------------|--------------------|------------|----|----|----|-------------|-----------|-----|-----|------------|------------|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Anjaw | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Changlang | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Dibang Valley | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| East Kameng | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| East Siang | NR | NR | NR | NR | NR | VHR | NR | NR | NR | VHR | NR | NR | NR |
| Kurung Kumey | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Lohit | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Lower Dibang Valley | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Lower Subansiri | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Papum Pare | NR | NR | NR | NR | NR | VHR | HR | NR | NR | VHR | NR | NR | NR |
| Tawang | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Tirap | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Upper Siang | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Upper Subansiri | NR | NR | NR | NR | NR | VHR | NR | NR | NR | VHR | VHR | NR | NR |
| West Kameng | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| West Siang | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Assam

| Districts of Assam | Livestock Diseases | | | | | | | | | | | | |
|---------------------|--------------------|------------|-----|----|-----|-------------|-----|-----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Baksa | NR | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | VLR | NR | NR |
| Barpeta | NR | NR | NR | NR | NR | NR | NR | HR | NR | NR | VLR | NR | NR |
| Bongaigaon | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Cachar | NR | NR | VHR | NR | NR | VHR | VLR | VHR | NR | NR | NR | NR | NR |
| Chirang | NR | NR | VHR | NR | NR | VHR | NR | VHR | NR | NR | VLR | NR | NR |
| Darrang | NR | NR | NR | NR | VHR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Dhemaji | NR | NR | NR | NR | NR | VHR | NR | VLR | NR | NR | VLR | NR | NR |
| Dhubri | NR | NR | VHR | NR | NR | VHR | VLR | NR | VHR | NR | NR | NR | NR |
| Dibrugarh | NR | NR | VHR | NR | NR | HR | VLR | VLR | NR | NR | NR | NR | NR |
| Dima Hasao | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Goalpara | NR | NR | VHR | NR | NR | NR | VLR | NR | VHR | VHR | VLR | NR | NR |
| Golaghat | NR | NR | VHR | NR | NR | VHR | NR | MR | NR | NR | VHR | NR | NR |
| Hailakandi | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Jorhat | NR | NR | VHR | NR | NR | VHR | NR | NR | NR | NR | VHR | NR | NR |
| Kamrup | NR | NR | VHR | NR | NR | VHR | NR | NR | NR | NR | VLR | NR | NR |
| Kamrup Metropolitan | NR | NR | HR | NR | NR | VHR | NR | VHR | NR | NR | HR | NR | NR |
| Karbi Anglong | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | VLR | NR | NR |
| Karimganj | NR | NR | VHR | NR | NR | VHR | NR | NR | NR | NR | VLR | NR | NR |
| Kokrajhar | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR | VLR | NR | NR |
| Lakhimpur | NR | NR | NR | NR | NR | MR | NR | NR | NR | NR | VLR | NR | NR |
| Morigaon | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

Continued

| Districts of Assam | Livestock Diseases | | | | | | | | | | | | |
|--------------------|--------------------|------------|-----|----|-----|-------------|-----|-----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Nagaon | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Nalbari | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Sivasagar | NR | NR | NR | NR | VHR | VHR | VLR | NR | HR | NR | HR | NR | NR |
| Sonitpur | NR | VHR | VHR | NR | NR | NR | VLR | MR | VHR | NR | VHR | NR | NR |
| Tinsukia | NR | VHR | NR | NR | NR | NR | VLR | VLR | VHR | NR | NR | VHR | NR |
| Udalguri | NR | NR | NR | NR | NR | VHR | VLR | HR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR), Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Bihar

| Districts of Bihar | Livestock Diseases | | | | | | | | | | | | |
|--------------------|--------------------|------------|----|----|----|-------------|-----|-----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Araria | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Arwal | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Aurangabad | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Banka | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Begusarai | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bhagalpur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bhojpur | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | VLR | NR | VHR |
| Buxar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Darbhanga | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Gaya | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Gopalganj | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR |
| Jamui | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Jehanabad | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kaimur (Bhabua) | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Katihar | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Khagaria | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | VHR |
| Kishanganj | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR |
| Lakhisarai | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Madhepura | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Madhubani | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Munger | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Muzaffarpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

Continued

| Districts of Bihar | Livestock Diseases | | | | | | | | | | | | |
|--------------------|--------------------|------------|-----|-----|----|-------------|-----|-----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Nalanda | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Nawada | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Pashchim Champaran | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Patna | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | VHR | NR | NR |
| Purba Champaran | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Purnia | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Rohtas | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Saharsa | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Samastipur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Saran | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Sheikhpura | NR | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Sheohar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Sitamarhi | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Siwan | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Supaul | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Vaishali | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Chandigarh

| Districts of Chandigarh | Livestock Diseases | | | | | | | | | | | | |
|-------------------------|--------------------|------------|----|----|----|-------------|-----|----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Chandigarh | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (**HR**), Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Chhattisgarh

| Districts of Chhattisgarh | Livestock Diseases | | | | | | | | | | | | |
|---------------------------|--------------------|------------|----|-----|----|-------------|-----|----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Bastar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Bijapur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bilaspur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR |
| Dakshin Bastar Dantewada | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Dhamtari | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Durg | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Janjgir-champa | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR |
| Jashpur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kabeerdham | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Korba | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Koriya | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Mahasamund | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Narayanpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Raigarhh | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Raipur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Rajnandgaon | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Surguja | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Uttar Bastar Kanker | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Dadra and Nagar Haveli

| Districts of Dadra and Nagar Haveli | Livestock Diseases | | | | | | | | | | | | |
|-------------------------------------|--------------------|------------|----|----|----|-------------|-----|----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Dadra and Nagar Haveli | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Daman and Diu

| Districts of Daman and Diu | Livestock Diseases | | | | | | | | | | | | |
|-------------------------------|--------------------|------------|----|-----|----|-------------|-----|-----|-----|------------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Daman | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Diu | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR | VLR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR), Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Goa

| Districts of Goa | Livestock Diseases | | | | | | | | | | | | |
|------------------|--------------------|------------|----|----|----|-------------|-----|-----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| North Goa | NR | VHR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | VHR | NR |
| South Goa | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR), Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Gujarat

| Districts of Gujarat | Livestock Diseases | | | | | | | | | | | | |
|----------------------|--------------------|------------|----|-----|-----|-------------|-----|-----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Ahmadabad | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Amreli | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Anand | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Banas Kantha | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Bharuch | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bhavnagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Dohad | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Gandhinagar | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR |
| Jamnagar | NR | NR | NR | NR | NR | NR | VHR | VLR | NR | VHR | NR | NR | NR |
| Junagadh | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | VLR | NR | NR |
| Kachchh | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Kheda | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Mahesana | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Narmada | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Navsari | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Panch Mahals | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Patan | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Porbandar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Rajkot | NR | NR | NR | NR | VHR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Sabar Kantha | NR | NR | NR | NR | NR | NR | VLR | NR | HR | NR | NR | NR | NR |
| Surat | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR |
| Surendranagar | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Tapi | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| The Dangs | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Vadodara | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Valsad | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Haryana

| Districts of Haryana | Livestock Diseases | | | | | | | | | | | | |
|----------------------|--------------------|------------|----|-----|-----|-------------|-----|-----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Ambala | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR |
| Bhiwani | NR | NR | NR | NR | VHR | NR | NR | VLR | NR | NR | NR | NR | NR |
| Faridabad | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Fatehabad | NR | NR | NR | NR | NR | NR | VLR | NR | VHR | NR | NR | NR | NR |
| Gurgaon | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Hisar | NR | NR | NR | NR | VHR | NR | NR | VLR | VHR | NR | VHR | NR | NR |
| Jhajjar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Jind | NR | NR | NR | NR | NR | NR | VLR | VHR | NR | NR | NR | NR | NR |
| Kaithal | NR | NR | NR | NR | NR | NR | HR | HR | NR | NR | VLR | NR | NR |
| Karnal | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Kurukshetra | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Mahendragarh | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Mewat | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Palwal | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Panchkula | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Panipat | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | VHR | NR |
| Rewari | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Rohtak | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Sirsa | NR | NR | NR | NR | NR | NR | NR | VLR | VHR | NR | NR | NR | NR |
| Sonipat | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Yamunanagar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Himachal Pradesh

| Districts of Himachal Pradesh | Livestock Diseases | | | | | | | | | | | | |
|----------------------------------|--------------------|------------|----|-----|----|-------------|-----|-----|-----|------------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Bilaspur | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Chamba | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Hamirpur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kangra | NR | NR | NR | NR | NR | NR | MR | NR | NR | NR | NR | NR | NR |
| Kinnaur | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Kullu | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Lahul & Spiti | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Mandi | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Shimla | NR | NR | NR | VLR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Sirmaur | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR |
| Solan | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Una | NR | NR | NR | NR | NR | NR | HR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Jammu and Kashmir

| Districts of Jammu and Kashmir | Livestock Diseases | | | | | | | | | | | | |
|--------------------------------|--------------------|------------|----|-----|----|-------------|-----|----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Anantnag | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Badgam | NR | NR | NR | VLR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Bandipore | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Baramula | NR | NR | NR | VLR | NR | NR | NR | NR | NR | HR | NR | NR | NR |
| Doda | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Ganderbal | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Jammu | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR |
| Kargil | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kathua | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Kishtwar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kulgam | NR | NR | NR | VLR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Kupwara | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Leh(Ladakh) | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Pulwama | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Punch | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Rajouri | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Ramban | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Reasi | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Samba | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Shupiyan | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Srinagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Udhampur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Jharkhand

| Districts of Jharkhand | Livestock Diseases | | | | | | | | | | | | |
|------------------------|--------------------|------------|-----|-----|------------|-------------|------------|------------|------------|------------|------------|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Bokaro | NR | VHR | NR | NR | VHR | VHR | VHR | VLR | VHR | NR | VHR | VHR | VHR |
| Chatra | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | VHR |
| Deoghar | NR | VHR | NR | NR | VHR | VHR | VHR | VHR | VHR | NR | NR | VHR | VHR |
| Dhanbad | NR | VHR | NR | NR | NR | VHR | VHR | VLR | VHR | NR | NR | VHR | VHR |
| Dumka | NR | VHR | NR | NR | NR | VHR | VHR | VHR | VHR | NR | VHR | VHR | VHR |
| Garhwa | NR | VHR | NR | VLR | NR | VHR | NR | NR | NR | NR | NR | VHR | HR |
| Giridih | NR | VHR | NR | VLR | VHR | VHR | HR | NR | NR | NR | HR | VHR | VHR |
| Godda | NR | VHR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | HR | VHR |
| Gumla | NR | VHR | NR | NR | NR | VHR | HR | MR | NR | NR | MR | VHR | VHR |
| Hazaribagh | NR | VHR | NR | VLR | NR | VHR | VLR | MR | NR | NR | NR | VHR | VHR |
| Jamtara | NR | VHR | NR | NR | NR | VHR | HR | VLR | HR | NR | NR | NR | VHR |
| Khunti | NR | VHR | NR | VLR | NR | VHR | VHR | VHR | VHR | NR | VHR | VHR | VHR |
| Koderma | NR | NR | NR | NR | VHR | VHR | NR | NR | NR | NR | NR | VHR | NR |
| Latehar | NR | VHR | NR | NR | VHR | VHR | VLR | NR | NR | NR | NR | VHR | VHR |
| Lohardaga | NR | VHR | NR | NR | NR | VHR | MR | NR | VHR | NR | VHR | VHR | VHR |
| Pakur | NR | VHR | NR | NR | VHR | VHR | VLR | NR | NR | NR | NR | VHR | VHR |
| Palamu | NR | VHR | NR | NR | VHR | VHR | VLR | NR | NR | NR | NR | VHR | VHR |
| RashchimiSinghbhum | NR | VHR | NR | VLR | NR | VHR | VHR | MR | MR | NR | MR | VHR | VHR |
| Purbi Singhbhum | NR | VHR | NR | NR | NR | VHR | VLR | NR | NR | NR | NR | VHR | VHR |
| Ramgarh | NR | VHR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | VHR | VHR |
| Ranchi | NR | VHR | VHR | VLR | NR | VHR | MR | NR | VHR | NR | MR | VHR | VHR |
| Sahibganj | NR | VHR | VHR | VLR | VHR | VHR | VHR | VHR | VHR | VHR | VHR | VHR | VHR |
| Seraikela - Kharsawan | NR | HR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | VHR | HR |
| Simdega | NR | VHR | NR | VLR | NR | VHR | MR | VLR | NR | NR | MR | VHR | VHR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR), Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Karnataka

| Districts of Karnataka | Livestock Diseases | | | | | | | | | | | | |
|------------------------|--------------------|------------|------------|-----|------------|-------------|------------|-----------|------------|------------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Bagalkot | NR | NR | NR | VLR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Bangalore | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Bangalore Rural | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR |
| Belgaum | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bellary | VHR | NR | NR | LR | VHR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Bidar | NR | NR | NR | VLR | NR | NR | VLR | VLR | NR | MR | NR | NR | NR |
| Bijapur | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Chamarajanagar | MR | NR | VHR | VLR | NR | NR | HR | NR | NR | NR | NR | NR | NR |
| Chikkaballapura | NR | NR | NR | LR | VHR | NR | VHR | NR | NR | MR | NR | NR | NR |
| Chikmagalur | NR | NR | MR | VLR | NR | NR | VHR | VLR | NR | NR | NR | NR | NR |
| Chitradurga | NR | NR | NR | LR | NR | NR | VLR | HR | VHR | HR | NR | NR | NR |
| Dakshina Kannada | NR | NR | NR | NR | NR | NR | MR | VLR | NR | NR | VLR | NR | NR |
| Davanagere | VHR | NR | VHR | VLR | NR | NR | NR | HR | NR | VHR | NR | NR | NR |
| Dharwad | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Gadag | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | MR | NR | NR | NR |

Continued

| Districts of Karnataka | Livestock Diseases | | | | | | | | | | | | |
|------------------------|--------------------|------------|------------|-----|------------|-------------|------------|------------|------------|------------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Gulbarga | VHR | NR | NR | VLR | NR | NR | VHR | VLR | NR | NR | NR | NR | NR |
| Hassan | NR | NR | VHR | NR | NR | NR | HR | NR | NR | NR | NR | NR | NR |
| Haveri | HR | NR | NR | NR | NR | NR | VLR | NR | NR | VHR | NR | NR | NR |
| Kodagu | NR | NR | MR | VLR | NR | NR | HR | NR | MR | NR | VLR | NR | NR |
| Kolar | NR | NR | NR | VLR | NR | NR | VHR | VLR | VHR | VHR | NR | NR | NR |
| Koppal | VHR | NR | NR | VLR | VHR | NR | VLR | VLR | NR | VHR | NR | NR | NR |
| Mandya | NR | NR | NR | NR | VHR | NR | VHR | VLR | NR | HR | NR | NR | NR |
| Mysore | MR | NR | VHR | VLR | NR | NR | VHR | VLR | NR | HR | NR | NR | NR |
| Raichur | HR | NR | NR | VLR | NR | NR | VLR | VLR | NR | VHR | NR | NR | NR |
| Ramanagara | NR | NR | NR | NR | NR | NR | VHR | VLR | NR | HR | NR | NR | NR |
| Shimoga | NR | NR | VHR | VLR | NR | NR | HR | NR | HR | VHR | NR | NR | NR |
| Tumkur | VHR | NR | HR | VLR | VHR | NR | MR | VHR | NR | VHR | NR | NR | NR |
| Udupi | NR | NR | NR | NR | NR | NR | MR | NR | NR | NR | NR | NR | NR |
| Uttara Kannada | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Yadgir | NR | NR | NR | VLR | NR | NR | VLR | VLR | VHR | VHR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Kerala

| Districts of Kerala | Livestock Diseases | | | | | | | | | | | | |
|---------------------|--------------------|------------|----|-----|----|-------------|-----|------------|------------|---------|------------|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Alappuzha | NR | NR | NR | NR | NR | NR | VHR | NR | VHR | NR | NR | NR | NR |
| Ernakulum | NR | VHR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR |
| Idukki | VHR | NR | NR | NR | NR | NR | HR | NR | NR | NR | NR | VHR | NR |
| Kannur | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR |
| Kasaragod | NR | NR | NR | VLR | NR | NR | MR | NR | NR | NR | VHR | NR | NR |
| Kollam | NR | NR | NR | NR | NR | NR | VHR | VHR | NR | NR | NR | NR | NR |
| Kottayam | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR |
| Kozhikode | VHR | NR | NR | NR | NR | NR | HR | NR | VHR | NR | NR | NR | NR |
| Malappuram | NR | NR | NR | NR | NR | NR | VHR | HR | NR | NR | NR | NR | NR |
| Palakkad | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | VHR | NR |
| Pathanamthitta | NR | NR | NR | NR | NR | NR | VHR | VLR | NR | NR | NR | NR | NR |
| Thiruvananthapuram | NR | NR | NR | NR | NR | NR | VHR | MR | HR | NR | HR | NR | NR |
| Thrissur | VHR | NR | NR | NR | NR | NR | VHR | VLR | NR | NR | NR | NR | NR |
| Wayanad | NR | HR | NR | NR | NR | NR | VHR | NR | VHR | NR | VLR | HR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Lakshadweep

| Districts of Lakshadweep | Livestock Diseases | | | | | | | | | | | | |
|-----------------------------|--------------------|------------|----|----|----|-------------|-----|----|-----|------------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Lakshadweep | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Madhya Pradesh

| Districts of Madhya Pradesh | Livestock Diseases | | | | | | | | | | | | |
|-----------------------------|--------------------|------------|----|-----|----|-------------|-----|-----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Alirajpur | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Anuppur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Ashoknagar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Balaghat | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Barwani | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Betul | NR | NR | NR | NR | NR | NR | HR | NR | NR | NR | NR | NR | NR |
| Bhind | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bhopal | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Burhanpur | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Chhatarpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Chhindwara | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Damoh | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Datia | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Dewas | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Dhar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Dindori | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| East Nimar | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Guna | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Gwalior | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Harda | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Hoshangabad | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Indore | NR | NR | NR | NR | NR | NR | VLR | VHR | NR | NR | NR | NR | NR |
| Jabalpur | NR | NR | NR | NR | NR | NR | VLR | HR | NR | NR | NR | NR | NR |
| Jhabua | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Katni | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

Continued

| Districts of Madhya Pradesh | Livestock Diseases | | | | | | | | | | | | |
|-----------------------------|--------------------|------------|----|-----|----|-------------|-----|----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Khargone (West Nimar) | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Mandla | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Mandsaur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Morena | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Narsimhapur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Neemuch | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Panna | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Raisen | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Rajgarh | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Ratlam | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Rewa | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Sagar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Satna | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Sehore | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Seoni | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Shahdol | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Shajapur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Sheopur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Shivpuri | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Sidhi | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Singrauli | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Tikamgarh | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Ujjain | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Umaria | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Vidisha | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR), Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Maharashtra

| Districts of Maharashtra | Livestock Diseases | | | | | | | | | | | | |
|--------------------------|--------------------|------------|-----|-----|----|-------------|-----|-----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Ahmadnagar | NR | NR | VHR | NR | NR | NR | VHR | VHR | VHR | NR | NR | VHR | NR |
| Akola | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Amravati | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Aurangabad | NR | NR | NR | NR | NR | NR | HR | NR | NR | NR | NR | NR | NR |
| Bhandara | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Bid | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Buldana | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Chandrapur | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Dhule | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Gadchiroli | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Gondiya | NR | NR | NR | VLR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Hingoli | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Jalgaon | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Jalna | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kolhapur | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Latur | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR | NR | NR | NR |
| Mumbai | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Mumbai Suburban | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Nagpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Nanded | NR | NR | NR | VLR | NR | NR | VLR | NR | VHR | NR | NR | NR | NR |
| Nandurbar | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Nashik | NR | NR | NR | NR | NR | NR | VHR | VHR | VHR | VHR | NR | VHR | NR |
| Osmanabad | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

Continued

| Districts of Maharashtra | Livestock Diseases | | | | | | | | | | | | |
|--------------------------|--------------------|------------|----|-----|-----|-------------|-----|----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Parbhani | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Pune | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Raigarh | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Ratnagiri | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Sangli | NR | NR | NR | NR | NR | NR | NR | NR | VHR | VHR | NR | VHR | NR |
| Satara | NR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR |
| Sindhudurg | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Solapur | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Thane | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Wardha | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Washim | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Yavatmal | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | VHR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Manipur

| Districts of Manipur | Livestock Diseases | | | | | | | | | | | | |
|----------------------|--------------------|------------|-----|----|-----|-------------|-----|-----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Bishnupur | NR | NR | VHR | NR | NR | VHR | NR | NR | NR | NR | NR | NR | NR |
| Chandel | NR | NR | VHR | NR | VHR | NR | NR | VLR | NR | NR | HR | NR | NR |
| Churachandpur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR |
| Imphal East | NR | NR | VHR | NR | NR | VHR | NR | NR | NR | NR | VLR | NR | NR |
| Imphal West | NR | NR | VHR | NR | NR | VHR | NR | VLR | NR | NR | VHR | NR | NR |
| Senapati | NR | NR | NR | NR | NR | VHR | NR | VLR | NR | HR | NR | NR | NR |
| Tamenglong | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | HR | NR | NR |
| Thoubal | NR | NR | NR | NR | NR | VHR | NR | NR | NR | VHR | HR | NR | NR |
| Ukhrul | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Meghalaya

| Districts of Meghalaya | Livestock Diseases | | | | | | | | | | | | |
|------------------------|--------------------|------------|------------|-----|----|-------------|------------|------------|-----|---------|------------|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| East Garo Hills | NR | NR | HR | NR | NR | NR | VLR | VHR | NR | NR | VHR | NR | NR |
| East Jaintia Hills | NR | NR | VHR | VLR | NR | NR | VLR | VLR | NR | NR | VHR | NR | NR |
| East Khasi Hills | NR | NR | VHR | NR | NR | NR | VHR | VLR | NR | NR | VHR | NR | NR |
| Jaintia Hills | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | MR | NR | NR |
| North Garo Hills | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR |
| Ribhoi | NR | NR | NR | NR | NR | NR | HR | NR | NR | NR | MR | NR | NR |
| South Garo Hills | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Southwest Garo Hills | NR | NR | VHR | NR | NR | NR | MR | VLR | NR | NR | NR | NR | NR |
| Southwest Khasi Hills | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | VLR | NR | NR |
| West Garo Hills | NR | NR | HR | NR | NR | NR | HR | VHR | NR | NR | VHR | NR | NR |
| West Khasi Hills | NR | NR | VHR | NR | NR | NR | VLR | VHR | NR | NR | LR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Mizoram

| Districts of Mizoram | Livestock Diseases | | | | | | | | | | | | |
|----------------------|--------------------|------------|----|----|----|-------------|-----|-----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Aizawl | MR | NR | NR | NR | NR | NR | HR | VLR | NR | NR | VHR | NR | NR |
| Champhai | NR | NR | NR | NR | NR | NR | VLR | VHR | NR | NR | VHR | NR | NR |
| Kolasib | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | VHR | NR | NR |
| Lawngtlai | NR | NR | NR | NR | NR | NR | VLR | NR | NR | VHR | VHR | NR | NR |
| Lunglei | NR | NR | NR | NR | NR | NR | MR | VLR | NR | NR | VHR | NR | NR |
| Mamit | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Saiha | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | VHR | VHR | NR | NR |
| Serchhip | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | VHR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Nagaland

| Districts of Nagaland | Livestock Diseases | | | | | | | | | | | | |
|-----------------------|--------------------|------------|----|----|----|-------------|-----|----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Dimapur | NR | NR | NR | NR | NR | NR | MR | NR | NR | NR | MR | NR | NR |
| Kiphire | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Kohima | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Longleng | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR |
| Mokokchung | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Mon | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Peren | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Phek | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Tuensang | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR |
| Wokha | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR |
| Zunheboto | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | HR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: NCT of Delhi

| Districts of NCT of Delhi | Livestock Diseases | | | | | | | | | | | | |
|---------------------------|--------------------|------------|----|-----|----|-------------|-----|----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Central | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| East | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| New Delhi | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| North | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| North East | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| North West | NR | NR | NR | NR | NR | NR | HR | NR | NR | NR | NR | NR | NR |
| South | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| South West | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| West | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Odisha

| Districts of Odisha | Livestock Diseases | | | | | | | | | | | | |
|---------------------|--------------------|------------|------------|-----|----|-------------|-----|------------|------------|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Anugul | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Balangir | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR |
| Baleshwar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Bargarh | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Baudh | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bhadrak | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Cuttack | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Debagarh | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Dhenkanal | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Gajapati | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Ganjam | NR | NR | NR | VLR | NR | NR | VLR | VHR | VHR | NR | NR | NR | NR |
| Jagatsinghpur | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Jajapur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Jharsuguda | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kalahandi | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Kandhamal | NR | NR | NR | VLR | NR | NR | VLR | NR | VHR | NR | NR | NR | NR |
| Kendrapara | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | VHR | NR |
| Kendujhar | NR | NR | VHR | NR | NR | NR | MR | VLR | NR | NR | NR | NR | NR |
| Khordha | NR | NR | NR | NR | NR | NR | MR | VLR | NR | NR | NR | NR | NR |
| Koraput | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Malkangiri | NR | NR | NR | VLR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |

Continued

| Districts of Odisha | Livestock Diseases | | | | | | | | | | | | |
|---------------------|--------------------|------------|-----|-----|----|-------------|-----|----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Mayurbhanj | NR | NR | NR | VLR | NR | NR | HR | NR | HR | NR | NR | MR | NR |
| Nabarangapur | NR | NR | VHR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR |
| Nayagarh | NR | NR | NR | VLR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Nuapada | NR | NR | VHR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Puri | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Rayagada | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Sambalpur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Subarnapur | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Sundargarh | NR | NR | NR | NR | NR | NR | MR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Puducherry

| Districts of Puducherry | Livestock Diseases | | | | | | | | | | | | |
|-------------------------|--------------------|------------|----|-----|----|-------------|-----|-----|-----|------------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Karaikal | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Mahe | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Puducherry | NR | VHR | NR | VLR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Yanam | NR | NR | NR | VLR | NR | VHR | VLR | VLR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Punjab

| Districts of Punjab | Livestock Diseases | | | | | | | | | | | | |
|----------------------------|--------------------|------------|----|----|----|-------------|-----|----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Amritsar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Barnala | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bathinda | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Faridkot | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Fatehgarh Sahib | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR |
| Firozpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VHR | NR | NR |
| Gurdaspur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Hoshiarpur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Jalandhar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kapurthala | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Ludhiana | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Mansa | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Moga | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Muktsar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Patiala | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | MR | NR | NR |
| Rupnagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Sahibzada Ajit Singh Nagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Sangrur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Shahid Bhagat Singh Nagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Tarn Taran | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Rajasthan

| Districts of Rajasthan | Livestock Diseases | | | | | | | | | | | | |
|------------------------|--------------------|------------|----|-----|----|-------------|------------|------------|------------|---------|------------|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Ajmer | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Alwar | NR | VHR | NR | NR | NR | NR | MR | HR | NR | NR | NR | NR | NR |
| Banswara | NR | NR | NR | NR | NR | VHR | VLR | NR | NR | NR | NR | NR | NR |
| Baran | NR | NR | NR | NR | NR | NR | VLR | VLR | VHR | NR | NR | NR | NR |
| Barmer | NR | VHR | NR | VLR | NR | NR | NR | VLR | NR | NR | NR | NR | NR |
| Bharatpur | NR | NR | NR | NR | NR | NR | HR | NR | NR | NR | NR | NR | NR |
| Bhilwara | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Bikaner | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bundi | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Chittaurgarh | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Churu | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Dausa | NR | NR | NR | VLR | NR | NR | VLR | HR | NR | NR | NR | NR | NR |
| Dhaulpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Dungarpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Ganganagar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Hanumangarh | NR | NR | NR | NR | NR | NR | VLR | VHR | NR | NR | NR | NR | NR |
| Jaipur | NR | NR | NR | NR | NR | NR | VHR | VHR | VHR | NR | NR | NR | NR |
| Jaisalmer | NR | NR | NR | VLR | NR | NR | VLR | NR | VHR | NR | NR | NR | NR |
| Jalor | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Jhalawar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VHR | NR | NR |
| Jhunjhunun | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Jodhpur | NR | NR | NR | VLR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Karauli | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

Continued

| Districts of Rajasthan | Livestock Diseases | | | | | | | | | | | | |
|------------------------|--------------------|------------|----|-----|------------|-------------|-----|----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Kota | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Nagaur | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Pali | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Pratapgarh | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Rajsamand | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Sawai Madhopur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Sikar | NR | NR | NR | NR | VHR | NR | MR | NR | NR | NR | NR | NR | NR |
| Sirohi | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Tonk | NR | NR | NR | NR | NR | NR | VLR | NR | MR | NR | NR | NR | NR |
| Udaipur | NR | NR | NR | NR | VHR | NR | VLR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Sikkim

| Districts of Sikkim | Livestock Diseases | | | | | | | | | | | | |
|---------------------|--------------------|------------|----|----|----|-------------|-----|-----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| East District | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | VHR | NR |
| North District | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR |
| South District | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| West District | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Tamil Nadu

| Districts of Tamil Nadu | Livestock Disease | | | | | | | | | | | | |
|-------------------------|-------------------|------------|-----|-----|----|-------------|-----|----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Ariyalur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Chennai | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Coimbatore | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Cuddalore | NR | NR | NR | NR | NR | NR | VLR | NR | VHR | NR | NR | NR | NR |
| Dharmapuri | VHR | NR | VHR | NR | NR | NR | VLR | NR | NR | VHR | NR | NR | NR |
| Dindigul | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Erode | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kancheepuram | VHR | NR | VHR | VLR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Kanniyakumari | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Karur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Krishnagiri | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Madurai | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Nagapattinam | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Namakkal | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Perambalur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Pudukkottai | VHR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Ramanathapuram | NR | NR | NR | VLR | NR | NR | NR | NR | NR | VHR | NR | NR | NR |
| Salem | NR | NR | NR | VLR | NR | NR | VHR | NR | NR | NR | NR | NR | NR |
| Sivaganga | NR | NR | NR | VLR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Thanjavur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| The Nilgiris | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Theni | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Thiruvallur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Thiruvarur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Thoothukkudi | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

Continued

| Districts of Tamil Nadu | Livestock Disease | | | | | | | | | | | | |
|-------------------------|-------------------|------------|----|-----|----|-------------|-----|------------|-----|------------|-----------|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Tiruchirappalli | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Tirunelveli | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | VHR | NR | NR | NR |
| Tiruppur | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Tiruvannamalai | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Vellore | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Viluppuram | NR | NR | NR | NR | NR | NR | VLR | VHR | NR | NR | HR | NR | NR |
| Virudhunagar | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Telangana

| Districts of Telangana | Livestock Diseases | | | | | | | | | | | | |
|---------------------------|--------------------|------------|----|-----|-----|-------------|-----|-----|-----|------------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Adilabad | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Hyderabad | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR | NR | NR | NR |
| Karimnagar | NR | NR | NR | MR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Khammam | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Mahbubnagar | NR | NR | NR | LR | VHR | NR | NR | VLR | NR | NR | NR | NR | NR |
| Medak | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Nalgonda | NR | NR | NR | VLR | NR | NR | NR | VLR | VHR | NR | NR | NR | NR |
| Nizamabad | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Rangareddy | NR | NR | NR | LR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Warangal | NR | NR | NR | VLR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease forewarning for March, 2022: Tripura

| Districts of Tripura | Livestock Disease | | | | | | | | | | | | |
|----------------------|-------------------|------------|------------|----|----|-------------|------------|------------|-----|------------|------------|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Dhalai | NR | NR | VHR | NR | NR | NR | VLR | VLR | NR | VHR | NR | NR | NR |
| North Tripura | NR | NR | NR | NR | NR | VHR | HR | VHR | NR | NR | VHR | NR | NR |
| South Tripura | NR | NR | VHR | NR | NR | VHR | VLR | HR | NR | VHR | VHR | NR | NR |
| West Tripura | VHR | NR | VHR | NR | NR | VHR | VHR | VHR | NR | VHR | VHR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Uttar Pradesh

| Districts of Uttar Pradesh | Livestock Disease | | | | | | | | | | | | |
|----------------------------|-------------------|------------|----|----|----|-------------|------------|----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Agra | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Aligarh | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | VLR | NR | NR |
| Allahabad | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | VHR | VHR |
| Ambedkar Nagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Amethi | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Auraiya | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Azamgarh | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Baghpat | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bahraich | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | VHR |
| Ballia | NR | VHR | NR | NR | NR | VHR | VLR | NR | NR | NR | VLR | VHR | VHR |
| Balrampur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Banda | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | VHR | VHR |
| Bara Banki | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bareilly | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | VHR |
| Basti | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bijnor | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR |
| Budaun | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR |
| Bulandshahr | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | VHR | VHR |
| Chandauli | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR |
| Chitrakoot | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | VHR |
| Deoria | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Etah | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | VHR |
| Etawah | NR | VHR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR | NR |
| Faizabad | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Farrukhabad | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

Continued

| Districts of Uttar Pradesh | Livestock Disease | | | | | | | | | | | | |
|----------------------------|-------------------|------------|----|----|------------|-------------|-----|-----|------------|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Fatehpur | NR | VHR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | VHR | VHR |
| Firozabad | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Gautam Buddha Nagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Ghaziabad | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Ghazipur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Gonda | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | HR | VHR |
| Gorakhpur | NR | VHR | NR | NR | NR | NR | VLR | NR | VHR | NR | NR | VHR | VHR |
| Hamirpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | MR | NR |
| Hapur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Hardoi | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | VHR |
| Jalaun | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | VHR |
| Jaunpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | HR |
| Jhansi | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Jyotiba Phule Nagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kannauj | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kanpur Dehat | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Kanpur Nagar | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | VHR |
| Kanshiram Nagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kaushambi | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | VHR | VHR |
| Kheri | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Kushinagar | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | VHR |
| Lalitpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Lucknow | NR | NR | NR | NR | VHR | VHR | NR | NR | NR | NR | NR | NR | VHR |
| MahaJanuarya Nagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Mahoba | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

Continued

| Districts of Uttar Pradesh | Livestock Disease | | | | | | | | | | | | |
|----------------------------|-------------------|------------|----|----|----|-------------|-----|-----|-----|---------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Mahrajganj | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR | VHR |
| Mainpuri | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Mathura | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | VHR | VHR |
| Mau | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Meerut | NR | VHR | NR | NR | NR | VHR | VHR | NR | NR | NR | NR | HR | VHR |
| Mirzapur | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | HR | VHR |
| Moradabad | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Muzaffarnagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Pilibhit | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Pratapgarh | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | HR |
| Rae Bareli | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | VHR | VHR |
| Rampur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VLR | NR | NR |
| Saharanpur | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR |
| Sambhal | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Sant Kabir Nagar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Sant Ravidas Nagar | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | NR | NR | NR |
| Shahjahanpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Shamli | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Shrawasti | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Siddharthnagar | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Sitapur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | VHR |
| Sonbhadra | NR | VHR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | VHR |
| Sultanpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Unnao | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Varanasi | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: Uttarakhand

| Districts of Uttarakhand | Livestock Disease | | | | | | | | | | | | |
|--------------------------|-------------------|------------|----|----|----|-------------|-----|-----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Almora | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Bageshwar | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Chamoli | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Champawat | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Dehradun | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Garhwal | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Hardwar | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | NR | NR | NR |
| Nainital | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Pithoragarh | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Rudraprayag | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Tehri Garhwal | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Udham Singh Nagar | NR | NR | NR | NR | NR | NR | VLR | VLR | NR | NR | MR | NR | NR |
| Uttarkashi | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

If vaccinated, please ignore the disease forecast.

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

District wise Livestock Disease Risk Forewarning for March, 2022: West Bengal

| Districts of West Bengal | Livestock Disease | | | | | | | | | | | | |
|----------------------------|-------------------|------------|------------|-----|----|-------------|------------|------------|------------|------------|-----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Bankura | NR | NR | VHR | NR | NR | NR | VLR | NR | VHR | NR | NR | HR | NR |
| Bardhaman | NR | NR | NR | NR | NR | NR | MR | HR | VHR | NR | NR | VHR | NR |
| Birbhum | NR | VHR | NR | VLR | NR | NR | VLR | NR | VHR | NR | NR | NR | VHR |
| Dakshin Dinajpur | NR | NR | NR | NR | NR | NR | MR | NR | NR | VHR | NR | NR | NR |
| Darjiling | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Haora | NR | VHR | VHR | NR | NR | NR | VLR | VHR | VHR | VHR | NR | VHR | NR |
| Hugli | NR | NR | VHR | NR | NR | NR | VHR | NR | HR | NR | NR | VHR | NR |
| Jalpaiguri | NR | NR | NR | NR | NR | VHR | VLR | NR | MR | NR | NR | NR | NR |
| Koch Bihar | NR | VHR | VHR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |
| Kolkata | NR | NR | MR | VLR | NR | NR | VLR | VLR | NR | NR | VLR | NR | NR |
| Maldah | NR | MR | NR | NR | NR | NR | VLR | VHR | NR | NR | NR | NR | NR |
| Murshidabad | NR | NR | NR | NR | NR | NR | VLR | NR | VHR | NR | NR | NR | NR |
| Nadia | NR | NR | NR | NR | NR | NR | MR | NR | HR | NR | NR | VHR | NR |
| North Twenty-Four Parganas | NR | NR | NR | NR | NR | NR | VLR | NR | NR | VHR | NR | NR | NR |
| Paschim Medinipur | NR | NR | HR | NR | NR | NR | VHR | MR | VHR | NR | NR | HR | NR |
| Purba Medinipur | NR | NR | VHR | NR | NR | NR | VHR | NR | NR | NR | NR | VHR | NR |
| Puruliya | NR | NR | NR | NR | NR | NR | VLR | VHR | VHR | NR | NR | NR | NR |
| South Twenty Four Parganas | NR | NR | VHR | NR | NR | NR | VLR | NR | NR | VHR | NR | NR | NR |
| Uttar Dinajpur | NR | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR |

*No risk/No data available (NR), Very low risk (VLR), Low risk (LR), Moderate risk (MR), High risk (HR),Very high risk (VHR)

II) Glimpse about the risk of predicted diseases:

The Livestock disease risk obtained based on the Machine Learning algorithm were further categorized into risk events using High Risk and Very High Risk.

1. Andaman and Nicobar

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|--------|--|---------------------------------------|------------------|
| 1. | Nicobars, North & Middle Andaman and South Andaman | Three | Fasciolosis |
| 2. | South Andaman | One | Sheep & Goat pox |

2. Andhra Pradesh

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|--------|-----------------------------|---------------------------------------|----------------------------|
| 1. | Anantapur, Kurnool, Y.S.R | Three | Anthrax |
| 2. | East Godavari | One | Black Quarter |
| 3. | Anantapur | One | Haemorrhagic Septicaemia |
| 4. | Sri Potti Sriramulu Nellore | One | Peste des Petits Ruminants |
| 5. | Sri Potti Sriramulu Nellore | One | Sheep & Goat pox |

3. Arunachal Pradesh

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|--------|--|---------------------------------------|------------------------|
| 1. | East Siang, Papum Pare, Upper Subansiri and West Siang | Four | Fasciolosis |
| 2. | Papum Pare | One | Foot and Mouth Disease |
| 3. | Papum Pare, Tawang and Upper Subansiri | Three | Sheep & Goat pox |
| 4. | East Siang and Upper Subansiri | Two | Swine Fever |

4. Assam

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|---|--|----------------------------|
| 1. | Sonitpur and Tinsukia | Two | Babesiosis |
| 2. | Baksa, Cachar, Chirang, Dhubri, Dibrugarh, Goalpara, Golaghat, Jorhat, Kamrup, Kamrup Metropolitan, Karimganj and Sonitpur | Twelve | Black Quarter |
| 3. | Darrang and Sivasagar | Two | Enterotoxaemia |
| 4. | Cachar, Chirang, Dhemaji, Dhubri, Dibrugarh, Golaghat, Jorhat, Kamrup, Kamrup Metropolitan, Karimganj, Sivasagar and Udalguri | Twelve | Fasciolosis |
| 5. | Barpeta, Cachar, Chirang, Kamrup Metropolitan and Udalguri | Five | Haemorrhagic Septicaemia |
| 6. | Darrang, Dhubri, Goalpara, Sivasagar, Sonitpur and Tinsukia | Six | Peste des Petits Ruminants |
| 7. | Goalpara | One | Sheep & Goat pox |
| 8. | Golaghat, Jorhat, Kamrup Metropolitan, Sivasagar and Sonitpur | Five | Swine Fever |
| 9. | Tinsukia | One | Theileriosis |

5. Bihar

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--------------------------------|--|---------------------|
| 1. | Jamui and Khagaria | Two | Babesiosis |
| 2. | Sheikhpura | One | Black Quarter |
| 3. | Patna | One | Swine Fever |
| 4. | Bhojpur and Khagaria | Two | Trypanosomosis |

6. Chhattisgarh

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--------------------------------|--|---------------------|
| 1. | Koriya | One | Anthrax |

7. Goa

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--------------------------------|--|---------------------|
| 1. | North Goa | One | Babesiosis |
| 2. | North Goa | One | Theileriosis |

8. Gujarat

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--------------------------------|--|----------------------------|
| 1. | Rajkot | One | Enterotoxaemia |
| 2. | Gandhinagar and Jamnagar | Two | Foot and Mouth Disease |
| 3. | Sabar Kantha and Valsad | Two | Peste des Petits Ruminants |
| 4. | Jamnagar | One | Sheep & Goat pox |

9. Haryana

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--------------------------------|--|----------------------------|
| 1. | Bhiwani and Hisar | Two | Enterotoxaemia |
| 2. | Kaithal | One | Foot and Mouth Disease |
| 3. | Kaithal and Jind | Two | Haemorrhagic Septicaemia |
| 4. | Sirsa | One | Peste des Petits Ruminants |
| 5. | Hisar | One | Swine Fever |
| 6. | Panipat | One | Theileriosis |
| 7. | Ambala | One | Trypanosomiasis |

10. Himachal Pradesh

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--------------------------------|--|----------------------------|
| 1. | Sirmaur and Una | Two | Foot and Mouth Disease |
| 2. | Bilaspur | One | Peste des Petits Ruminants |
| 3. | Kinnaur | One | Sheep & Goat pox |

11. Jammu & Kashmir

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--|--|---------------------|
| 1. | Anantnag, Badgam, Bandipore, Baramula, Kulgam, Ganderbal, Kulgam, Kupwara, Shupiyan and Srinagar | Ten | Sheep & Goat pox |


12. Jharkhand

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|---|--|------------------------|
| 1. | All twenty-three districts except Koderma | Twenty-three | Babesiosis |
| 2. | Ranchi, and Sahibganj | Two | Black Quarter |
| 3. | Bokaro, Deoghar, Giridih, Koderma, Latehar, Pakur, Palamu, and Sahibganj | Eight | Enterotoxaemia |
| 4. | All twenty-two districts except Deoghar and Ramghar | Twenty-two | Fasciolosis |
| 5. | Bokaro, Deoghar, Dhanbad, Dumka, Giridih, Gumla, Jamtara, Khunti, Pashchimi Singhbhum and Sahibganj | Ten | Foot and Mouth Disease |




| | | | |
|-----|---|--------------|----------------------------|
| 6. | Deoghar, Dumka, Khunti and Sahibganj | Four | Haemorrhagic Septicaemia |
| 7. | Bokaro, Deoghar, Dhanbad, Dumka, Jamtara, Khunti, Lohardaga, Ranchi and Sahibganj | Nine | Peste des Petits Ruminants |
| 8. | Khunti and Sahibganj | Two | Sheep & Goat pox |
| 9. | Bokaro, Dumka, Giridih, Lohardaga and Sahibganj | Five | Swine Fever |
| 10. | All twenty-three districts except Jamtara | Twenty-three | Theileriosis |
| 11. | All twenty-three districts except Koderma | Twenty-three | Trypanosomosis |

13. Karnataka



| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|--------|---|---------------------------------------|----------------------------|
| 1. | Bellary, Davanagere, Gulbarga, Haveri, Koppal, Raichur and Tumkur | Seven | Anthrax |
| 2. | Chamarajanagar, Davanagere, Hassan, Mysore, Shimoga and Tumkur | Six | Black Quarter |
| 3. | Bellary, Chikkaballapura, Koppal, Mandya, and Tumkur | Five | Enterotoxaemia |
| 4. | Bangalore Ruler, Chamarajanagar, Chikkaballapura, Chikmagalur, Gadag, Gulbarga, Kodagu, Kolar, Mandya, Mysore, Ramanagara and Shimoga | Twelve | Foot and Mouth Disease |
| 5. | Chitradurga, Davanagere and Tumkuru | Three | Haemorrhagic Septicaemia |
| 6. | Chitradurga, Kolar, Shimoga and Yadgir | Four | Peste des Petits Ruminants |
| 7. | Bagalkot, Chitradurga, Davanagere, Haveri, Kolar, Koppal, Mandya, Mysore, Raichur, Ramanagara, Shimoga, Tumkur and Yadgir | Thirteen | Sheep & Goat pox |

14. Kerala

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--|--|----------------------------|
| 1. | Idukki, Kozhikode and Thrissur | Three | Anthrax |
| 2. | Ernakulam and Wayanad | Two | Babesiosis |
| 3. | All Fourteen Districts except Kasaragod | Thirteen | Foot and Mouth Disease |
| 4. | Kollam and Malappuram | Two | Haemorrhagic Septicaemia |
| 5. | Alappuzha, Kozhikode, Thiruvananthapuram and Wayanad | Four | Peste des Petits Ruminants |
| 7. | Kasaragod and Thiruvananthapuram | Two | Swine Fever |
| 8. | Idukki, Palakkad, and Wayanad | Three | Theileriosis |

15. Madhya Pradesh

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--------------------------------|--|--------------------------|
| 1. | Betul | One | Foot and Mouth Disease |
| 2. | Indore and Jabalpur | Two | Haemorrhagic Septicaemia |

16. Maharashtra

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|---|--|----------------------------|
| 1. | Ahmadnagar | One | Black Quarter |
| 2. | Satara | One | Enterotoxaemia |
| 3. | Aurangabad, Aurangabad and Nashik | Three | Foot and Mouth Disease |
| 4. | Ahmadnagar and Nashik | Two | Haemorrhagic Septicaemia |
| 5. | Ahmadnagar, Dhule, Nanded, Nandurbar, Nashik and Sangli | Six | Peste des Petits Ruminants |
| 6. | Nashik, Sangli and Yavatmal | Three | Sheep & Goat pox |

| | | | |
|----|-------------------------------|-------|--------------|
| 7. | Ahmadnagar, Nashik and Sangli | Three | Theileriosis |
|----|-------------------------------|-------|--------------|

17. Manipur

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|--------|---|---------------------------------------|------------------|
| 1. | Bishnupur, Chandel, Imphal East and Imphal West | Four | Black Quarter |
| 2. | Chandel | One | Enterotoxaemia |
| 3. | Bishnupur, Imphal East, Imphal West, Senapati and Thoubal | Five | Fasciolosis |
| 4. | Senapati and Thoubal | Two | Sheep & Goat pox |
| 5. | Chandel, Imphal West, Tamenglong and Thoubal | Four | Swine Fever |

18. Meghalaya

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|--------|---|---------------------------------------|--------------------------|
| 1. | East Garo Hills, East Khasi Hills, East Khasi Hills, Southwest Garo Hills, West Garo Hills and West Khasi Hills | Six | Black Quarter |
| 2. | East Khasi Hills Jaintia Hills, Ribhoi and West Garo Hills | Four | Foot and Mouth Disease |
| 3. | East Garo Hills, West Garo Hills and West Khasi Hills | Three | Haemorrhagic Septicaemia |
| 4. | East Garo Hills, East Jaintia Hills, East Khasi Hills, North Garo Hills and West Garo Hills | Five | Swine fever |

19. Mizoram

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|---|--|--------------------------|
| 1. | Aizawl | One | Foot and Mouth Disease |
| 2. | Champhai | One | Haemorrhagic Septicaemia |
| 3. | Lawngtlai and Saiha | Two | Sheep & Goat pox |
| 4 | Aizawl, Champhai, Kolasib, Lawngtlai, Lunglei, Saiha and Serchhip | Seven | Swine Fever |

20. Nagaland



| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--------------------------------|--|---------------------|
| 1. | Zunheboto | One | Swine fever |

21. NCT of Delhi

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--------------------------------|--|------------------------|
| 1. | North West | One | Foot and Mouth Disease |

22. Odisha

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|-------------------------------------|--|--------------------------|
| 1. | Koraput | One | Anthrax |
| 2. | Gajapati | One | Babesiosis |
| 3. | Kendujhar, Nabarangapur and Nuapada | Three | Black Quarter |
| 4. | Mayurbhanj and Nabarangapur | Two | Foot and Mouth Disease |
| 5. | Ganjam | One | Haemorrhagic Septicaemia |

| | | | |
|----|--|------|----------------------------|
| 6. | Ganjam, Jagatsinghapur, Kandhamal, Mayurbhanj and Nayagarh | Five | Peste des Petits Ruminants |
| 7. | Kendrapara | One | Theileriosis |

23. Puducherry

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|--------|-------------------------|---------------------------------------|------------------|
| 1. | Karaikal and Puducherry | Two | Babesiosis |
| 2. | Yanam | One | Fasciolosis |
| 3. | Puducherry | One | Sheep & Goat pox |

24. Punjab

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|--------|-------------------------|---------------------------------------|--------------|
| 1. | Firozpur | One | Swine Fever |

25. Rajasthan

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|--------|--------------------------------------|---------------------------------------|----------------------------|
| 1. | Alwar and Barmer | Two | Babesiosis |
| 2. | Sikar and Udaipur | Two | Enterotoxaemia |
| 3. | Banswara | One | Fasciolosis |
| 4. | Bharatpur and Jaipur | Two | Foot and Mouth Disease |
| 5. | Alwar, Dausa, Hanumangarh and Jaipur | Four | Haemorrhagic Septicaemia |
| 6. | Baran, Jaipur, Jaisalmer and Jodhpur | Four | Peste des Petits Ruminants |
| 7. | Jhalawar | One | Swine Fever |

26. Sikkim

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|---------------------------------|--|---------------------|
| 1. | East District and West District | Two | Babesiosis |
| 2. | East District | One | Swine Fever |
| 3. | East District | One | Theileriosis |

27. Tamil Nadu

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--|--|----------------------------|
| 1. | Dharmapuri, Kancheepuram and Pudukkottai | Three | Anthrax |
| 2. | Dharmapuri and Kancheepuram | Two | Black Quarter |
| 3. | Salem | One | Foot and Mouth Disease |
| 4. | Viluppuram | One | Haemorrhagic Septicaemia |
| 5. | Cuddalore, Kancheepuram and Sivaganga | Three | Peste des Petits Ruminants |
| 6. | Dharmapuri, Ramanathapuram and Tirunelveli | Three | Sheep & Goat pox |
| 7. | Viluppuram | One | Swine Fever |

28. Telangana

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--------------------------------|--|----------------------------|
| 1. | Mahbubnagar | One | Enterotoxaemia |
| 2. | Nalgonda | One | Peste des Petits Ruminants |

29. Tripura

| Sl. No | Disease prone districts | Number of disease prone for districts | Disease Name |
|---------------|---|--|--------------------------|
| 1. | West Tripura | One | Anthrax |
| 2. | Dhalai, South Tripura and West Tripura | Three | Black Quarter |
| 3. | North Tripura, South Tripura and West Tripura | Three | Fasciolosis |
| 4. | North Tripura and West Tripura | Two | Foot and Mouth Disease |
| 5. | North Tripura, South Tripura and West Tripura | Three | Haemorrhagic Septicaemia |
| 6. | Dhalai, , South Tripura and West Tripura | Three | Sheep & Goat pox |
| 7. | North Tripura, South Tripura and West Tripura | Three | Swine Fever |


30. Uttar Pradesh

| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|---------------|--|--|----------------------------|
| 1. | Allahabad, Ballia, Banda, Bareilly, Bijnor, Chitrakoot, Etah, Etawah, Fatehpur, Gonda, Gorakhpur, Hardoi, Jalaun, Kanpur Nagar, Kaushambi, Kushinagar, Mathura, Meerut, Mirzapur, Saharanpur and Sonbhadra | Twenty-one | Babesiosis |
| 2. | Lucknow | One | Enterotoxaemia |
| 3. | Ballia, Etawah, Lucknow, Meerut and Sant Ravidas Nagar (Bhadohi) | Five | Fasciolosis |
| 4. | Budaun and Meerut | Two | Foot and Mouth Disease |
| 5. | Gorakhpur | One | Peste des Petits Ruminants |




| | | | |
|----|---|-------------|----------------|
| 6. | Allahabad, Ballia, Banda, Bareilly, Bulandshahr, Etah, Fatehpur, Gonda, Gorakhpur, Hardoi, Kanpur Nagar, Kaushambi, Kushinagar, Mahrajganj, Mathura, Meerut, Mirzapur and Rae Bareli, | Eighteen | Theileriosis |
| 7. | Allahabad, Bahraich, Ballia, Banda, Bareilly, Bijnor, Bulandshahr, Chandauli, Chitrakoot, Etah, Fatehpur, Gonda, Gorakhpur, Hardoi, Jalaun, Jaunpur, Kanpur Nagar, Kaushambi, Kushinagar, Lucknow, Mahrajganj, Mathura, Meerut, Mirzapur, Pratapgarh, Rae Bareli, Saharanpur, Sitapur and Sonbhadra | Twenty-nine | Trypanosomosis |

31. Uttarakhand




| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|--------|-------------------------|---------------------------------------|----------------------------|
| 1. | Rudraprayag | One | Peste des Petits Ruminants |

32. West Bengal



| Sl. No | Disease prone districts | Number of districts prone for disease | Disease Name |
|--------|--|---------------------------------------|------------------------|
| 1. | Birbhum, Haora and Koch Bihar | Three | Babesiosis |
| 2. | Bankura, Haora, Hugli, Koch Bihar, Paschim Medinipur, Purba Medinipur and South Twenty-Four Parganas | Seven | Black Quarter |
| 3. | Jalpaiguri | One | Fasciolosis |
| 4. | Hugli, Paschim Medinipur and Purba Medinipur | Three | Foot and Mouth Disease |

| | | | |
|----|--|-------|----------------------------|
| 4. | Barddhaman, Haora, Maldah and Puruliya | Four | Haemorrhagic Septicaemia |
| 5. | Bankura, Barddhaman, Birbhum, Haora, Hugli, Murshidabad, Nadia, Paschim Medinipur and Puruliya | Nine | Peste des Petits Ruminants |
| 6. | Dakshin Dinajpur, Haora, North Twenty Four Parganas and South Twenty Four Parganas | Four | Sheep & Goat pox |
| 7. | Bankura, Barddhaman, Haora, Hugli, Nadia, Paschim Medinipur and Purba Medinipur | Seven | Theileriosis |
| 8. | Birbhum | One | Trypanosomosis |



iii) Diseases, Species affected Clinical signs and its preventive measures.

| Sl No. | Disease | Species Affected | Clinical Signs | Preventive Measures |
|---------------|--------------------|--|---|---|
| 1 | Anthrax (AX) | Most of the mammals and ruminants are highly susceptible. Pigs and Horses are moderately susceptible. Carnivores are relatively resistant. | Convulsion and sudden death with oozing of blood from natural orifices such as rectum and nose prior to death. Occasionally oedema develops in the throat and shoulder over a period of one week before death. | Ring vaccination and reporting of the disease is advised. Vaccination to be done in consultation with the veterinarians and as decided by state animal husbandry authorities. Strict biosecurity measures may be followed. Carcass may be disposed by deep burying covered with lime powder. Contaminated area may be disinfected with 4% formalin or 10% caustic soda. Grazing area may be restricted. |
| 2 | Babesiosis (BA) | Cattle. Cross breeds are more susceptible. | High temperature, jaundice like symptoms, yellowish mucosal membrane of eye, rectum and coffee colour urine. | Periodical application of acaricides in and around the animal shed and on the animals. For therapeutic application, Diaminazine or Imidocarb can be useful. |
| 3. | Black Quarter (BQ) | Common disease of cattle and sheep, but occasionally goats and pigs also suffer from the disease. | High fever and lameness followed by swelling in the neck, shoulder, lumbar, gluteal and sacral regions. Skin over the affected area become dark and crepitate on palpation. Loss of feed intake, colic, lateral recumbency, dyspnoea and death. | Affected animals may be treated with suitable antibiotics. Vaccination to be done in consultation with the veterinarians and as decided by state animal husbandry authorities. Strict biosecurity measures may be followed. Grazing area may be restricted. Carcass may be disposed hygienically. |

| | | | | |
|----|---------------------|---|--|--|
| 4. | Bluetongue (BT) | Sheep are more susceptible than goats. | Fever, swelling of face, neck, eyelids respiratory distress, nasal discharge, Salivation, necrotic ulcers on tongue, dental pad, gum, lips hyperaemia of muzzle and may bleed at muco-cutaneous junction. Affected tongue may become swollen, cyanotic and purple blue in colour – ‘bluetongue’. | Vector control using insecticides and good water management. Vaccination of susceptible animals preferably in the month of May. Do not shear sheep during winter months. Restriction in animal movement, segregation of affected animals and symptomatic treatment. Strict biosecurity measures. |
| 5. | Enterotoxaemia (ET) | Common disease of sheep and goats especially among the young animals. | Dullness, opisthosomas, convulsions, coma and sudden death. Affected adult sheep, which survive for several days May show diarrhoea and staggering. | Affected animals may be treated with suitable antibiotics. Vaccination to be done in consultation with the veterinarians and as decided by State Animal Husbandry Authorities. Strict biosecurity measures may be followed. Carcass may be disposed hygienically. Grazing area to be restricted, stall fed, vitamins and probiotics may be provided. |
| 6. | Fasciolosis (FA) | Cattle, buffalo, sheep and goats. | Progressive anaemia, pale mucous membrane, sub-mandibular oedema (Bottle jaw), loss of appetite, weakness, isolated from flock while grazing, loss in production. | The animal should not be allowed to graze in water stagnant fields or submerged fodder should not be given directly to the animals. The submerged fodder can be processed through hay/silage preparation in order to destroy the metacercariae. The affected animals can be treated with Carbon tetrachloride/Rafoxanide/Nitroxynil/Niclofolan/Closantel/Oxyclozanide, |


| | | | | |
|----|-------------------------------|--|--|--|
| | | | | under the strict supervision of veterinarian. |
| 7. | Foot and Mouth Disease (FMD) | Cattle, buffalo, sheep, goats and pigs are often affected domesticated species, but the disease is more severe in cattle and pigs. | Fever, loss of feed intake, drop in milk production, drooling of saliva like ropey string, vesicles develop on the tongue, lips, gums, and palate and eventually rupture. Concurrent to oral lesions, vesicles also appear in inter digital skin and coronary band of the feet. The animal may open and close its mouth with a characteristic smacking sound. Sheep and goats may show lameness. In pigs, lesions may be seen on snout and also on the feet. | Regular vaccination and seromonitoring. Disinfection with sodium carbonate (4%) or 10% washing soda and strict biosecurity measures to be followed and animal movement may be controlled.  |
| 8. | Haemorrhagic septicaemia (HS) | Common disease for cattle and buffaloes, but can also occur among other species such as pigs, sheep, goats and many wild animals. | The disease starts with high fever, respiratory distress and haemorrhages maybe seen on the mucous membranes. There is lacrymation, nasal discharge, drop in milk production and anorexia. As the disease progress ear droops and the animals will be prostrated with cyanosis of mucous membranes. There may be oedema along the head, neck, thorax, vulva and anal areas. Sudden death occurs within few hours of clinical signs. | Affected animals may be treated with suitable antibiotics. Vaccination to be done in consultation with the veterinarians and as decided by state animal husbandry authorities. Strict biosecurity measures may be followed. Carcass may be disposed hygienically and stress factors may be reduced by following good animal husbandry practices. |

| | | | | |
|-----|----------------------------------|---|---|--|
| 9. | Peste des Petits Ruminants (PPR) | Goats and sheep are most affected domestic animals. | Fever, nasal and ocular discharge, respiratory distress, necrotic lesions in buccal mucosa, gum, dental pad, palate, tongue and diarrhoea. Animals may die because of dehydration and pneumonia. | Vaccination of susceptible animals of above 3 months old age. Restriction on animal movement, strict biosecurity measures and proper disposal of carcass. |
| 10. | Sheep and Goat pox (SGP) | Sheep and Goats | Respiratory distress and pock lesions over the non-hairy parts of body, more common in teat, udder, scrotum, head, neck, ear, perineum, inner aspect of thighs and under tail. | Vaccination of susceptible animals of above 3 months old age. Symptomatic treatment of affected animals. Restriction on animal movement, strict biosecurity measures and proper disposal of carcass. |
| 11. | Swine Fever (SF) | Pigs | Fever, Conjunctivitis, purplish discolouration of snout, ears, abdomen, inner side of the legs and staggering gait. | Vaccination of susceptible animals. Restriction on animal movement, strict biosecurity measures and proper disposal of carcass. |
| 12. | Theileriosis (TE) | Large Ruminants. Cross bred cattle are more vulnerable. | High temperature, yellowish eye, sometime eye maybe heavily swollen, icteric mucosal membrane of rectum, dark yellowish urine, sometime may reach to coffee colour. Antibiotic is of no use to check the fever. | Periodical application of acaricides in and around the animal shed and on the animals. Therapeutic treatment with Buparvaquone can be useful in both early and advanced stages of the infection. |


| | | | | |
|-----|---------------------|---|---|--|
| 13. | Trypanosomosis (TR) | Domestic and wild carnivores and herbivores including cattle, buffalo, horse, donkey, camel, dog and cats. Buffaloes are known as carriers. | Fluctuating high fever which is not responded by antibiotics, swollen lymph gland, chronic emaciation and weakness, loss of appetite, gradual loss of production. | The affected animal should be treated with Diaminazine compounds or chloride and sulphate salts of Quinapyramine. Periodical spray of insecticide in and around animal shed to remove the flies. |
|-----|---------------------|---|---|--|




iv) Risk Prediction - Livestock Disease Forewarning Maps




Risk Prediction of Babesiosis for the month of March 2022




Risk Prediction of Black quarter for the month of March 2022




Risk Prediction of Bluetongue for the month of March 2022




Risk Prediction of Enterotoxemia for the month of March 2022




Risk Prediction of Fascioliasis for the month of March 2022




Risk Prediction of Foot and mouth disease for the month of March 2022




Risk Prediction of Haemorrhagic septicaemia for the month of March 2022




Risk Prediction of Peste des petits ruminants for the month of March 2022




Risk Prediction of Sheep and Goat pox for the month of March 2022




Risk Prediction of Swine fever for the month of March 2022




Risk Prediction of Theileriosis for the month of March 2022




Risk Prediction of Trypanosomiasis for the month of March 2022



V. Forecasting of remote sensing and meteorological parameters till March, 2022(Ex. Tamil Nadu)





VI. SIGNIFICANT WEATHER PARAMETERS TABLE

| Disease Names | Significant Parameters |
|-----------------------------------|--|
| Anthrax | Precipitable Water, Surface Pressure |
| Babesiosis | Air Temperature, Precipitation, Surface Pressure, Minimum Temperature, Vapour Pressure, Dew Point Temperature, Wind Speed, NDVI and LST Night |
| Black quarter | Precipitable Water, Precipitation, Surface Pressure, Sea Level Pressure and Vapour Pressure |
| Bluetongue | Air Temperature, Cloud, Precipitable Water, Precipitation, Surface Pressure, Uwind, Vwind, Vapour Pressure, Elevation, NDVI and PET |
| Classical Swine Fever | Cloud, Precipitation, relative humidity Minimum Temperature, Vapour Pressure and Rainfall |
| Enterotoxaemia | Surface Pressure, NDVI and PET |
| Fasciolosis | Air Temperature, Precipitation, relative humidity, Temperature, Maximum Temperature, Vapour Pressure, Vwind, Rainfall, Soil Moisture, NDVI and EVI |
| Foot and Mouth Disease | Precipitable Water, Uwind, Vwind, wet_dry frequency, LST Night and EVI |
| Haemorrhagic Septicaemia | Cloud, Precipitation and Vwind |
| Peste des Petits Ruminants | Cloud, Precipitable Water, Surface Pressure, Sea Level Pressure, Maximum Temperature, Vwind and NDVI |
| Sheep & Goat pox | Cloud, Surface Pressure, Maximum Temperature, Vwind, DTP, NDVI and PET |
| Theileriosis | Air Temperature, Precipitation, Vapour Pressure, NDVI and LST Night |
| Trypanosomosis | Air Temperature, Precipitation, Surface Pressure, Vapour Pressure, NDVI and LST Night |

Table 7.1: Significant weather parameters govern the Livestock disease incidence (forecast).

9. Post prediction Validation

DIMAPUR | Publish Date: 4/14/2019 AH&VS TEAM VISITS AFFECTED AREAS UNDER MEDZIPHEMA,
Source: <http://www.nagalandpost.com>

Following reports of a good number of buffaloes dying in a recent outbreak of suspected Haemorrhagic septicaemia (HS), a team from Animal Husbandry and Veterinary Services (AH&VS) department visited the affected areas under Medziphema on April 12. (Haemorrhagic septicaemia is a contagious bacterial disease that affects cattle and water buffaloes with a high mortality rate in infected animals).

AH&VS, deputy director & principal investigator, AICRP-ADMAS, Dr S. Amenla Walling, in a press release reported that the team consisted of the department's director, Dr Temsumeren, along with additional director, Dr. Budhi Lama, and other officials from the department. The press release added that the area is prone to such kind of disease outbreaks and the department officials reminded villagers to cooperate with the department and vaccinate their animals against such outbreaks. The team told the villagers that even an outbreak can be contained more effectively if villagers report the matter on time to the nearest Veterinary Health Centre.


The villagers admitted in the meeting that they had not reported the recent outbreak to the department initially. The director appreciated the CVO Dimapur and his Rapid Response Team for their quick action after receiving information and for remaining stationed in the outbreak area to date. Free medicine was also distributed among the villagers. The department, through the press release also appealed to everyone to report such matters to the nearest Veterinary Health Centre (so that qualified staff may intervene quickly), instead of publicizing it in other ways. It stated that the department is prepared to extend services to any outbreak of diseases in animals to control such things.

The press release also pointed out that to control the recent outbreak, the department had to direct its officials to make their own transport arrangements to go to the affected areas because the State Election department did not consider an appeal to exempt the department's emergency duty vehicle from election duty.


Meanwhile, when contacted, Dr S. Amenla Walling told Nagaland Post that it is difficult to say if the disease has been fully contained since its free grazing season for the animals, but the department is doing its best under the circumstances.

NIVEDI PREDICITONS

| Districts of Nagaland | HS prediction for February 2019 | HS prediction for March 2019 | HS prediction for April 2019 |
|-----------------------|---------------------------------|------------------------------|------------------------------|
| Peren | VLR | VLR | VHR |
| Dimapur | VLR | NR | HR |
| Kohima | VLR | VLR | NR |
| Wokha | VLR | NR | VLR |



Risk Prediction of Haemorrhagic septicaemia for the month of April 2019



Note: Spatial and temporal neighbours

FMD POST PREDICTION

Published Date: 2021-04-14 14:06:16

Subject: PRO/SOAS> Foot & mouth disease - India: (Arunachal Pradesh) mithun

Archive Number: 20210414.8306219

FOOT & MOUTH DISEASE - INDIA: (ARUNACHAL PRADESH) MITHUN

A ProMED-mail post

<http://www.promedmail.org>

ProMED-mail is a program of the
International Society for Infectious Diseases
<http://www.isid.org>

Date: Tue 13 Apr 2021 6:32 AM IST

Source: The Sentinel [edited]

<https://www.sentinelassam.com/north-east-india-news/arunachal-news/many-mithuns-infected-with-fmd-foot-and-mouth-disease-533689>

A large number of mithuns [or gayal, a large domestic bovine] have been affected by FMD (foot-and-mouth disease) in various parts of Arunachal Pradesh, and a few have died, said official sources on Monday [13 Apr 2021].

Animal Husbandry & Veterinary Department Deputy director Dr Taba Heli, a top mithun expert in North Eastern region, reported that the disease has taken a severe form in the entire Siang belt, particularly in East Siang, West Siang, and Upper Siang districts.

Though the number of deaths is yet to be known, the disease has spread in scattered areas of Papum Pare district also. The Department has allocated district funds for procurement of medicines to take all possible steps to contain the disease, he said. Mithun deaths have been reported also from Itanagar and Jullang area.

Pointing out that mithuns are the pride of indigenous people in the state, Nyishi Elite Society president Bengia Tolum has urged the department to take all possible steps to save them.

--
Communicated by:

ProMED-SoAs from HealthMap Alerts


<promed-SoAs@promedmail.org>

District wise Livestock Disease Risk Forewarning for April 2021: Arunachal Pradesh

| Districts of Arunachal Pradesh | Livestock Diseases | | | | | | | | | | | | |
|--------------------------------|--------------------|------------|-----|----|----|--------------|-----|-----|-----|---------|-----|--------------|-----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fascioliasis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomiasis |
| Anjaw | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Changlang | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Dibang Valley | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| East Kameng | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| East Siang | NR | NR | NR | NR | NR | NR | VHR | NR | VLR | NR | VHR | NR | NR |
| Kurung Kumey | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Lohit | NR | VLR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Lower Dibang Valley | NR | NR | VLR | NR | NR | NR | VLR | VLR | VLR | NR | VHR | NR | NR |
| Lower Subansiri | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR | VLR | NR |
| Papum Pare | NR | NR | NR | NR | NR | NR | VHR | NR | VHR | NR | VHR | NR | NR |
| Tawang | NR | NR | NR | NR | NR | NR | HR | NR | NR | NR | NR | NR | NR |
| Tirap | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Upper Siang | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Upper Subansiri | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| West Kameng | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| West Siang | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

If vaccination has already been done please ignore the disease forecast for that disease.

Andaman and Nicobar Report June-2020



HIMACHAL PRADESH REPORT AUGUST-2020


NIVEDI Prediction

District wise Livestock Disease forewarning for July 2020: Himachal Pradesh


| Districts of Himachal pradesh | Livestock Diseases | | | | | | | | | | | | |
|-------------------------------|--------------------|------------|----|-----|----|-------------|-----|----|-----|---------|----|--------------|-----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomiasis |
| Bilaspur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Chamba | NR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR |
| Hamirpur | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kanera | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Kinnar | NR | NR | NR | VLR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Kullu | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Lahul & Spiti | NR | NR | NR | VLR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Mandi | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Shimla | NR | NR | NR | NR | NR | NR | NR | NR | VHR | NR | NR | NR | NR |
| Sirmaur | NR | NR | NR | NR | NR | NR | MR | NR | NR | NR | NR | NR | NR |
| Solan | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |
| Una | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR | NR |

If vaccination has already been done please ignore the disease forecast for that disease.

HIMACHAL PRADESH Risk Prediction of Sheep and Goat pox for the month of July 2020



HINACHAL PRADESH Report July-2020



| FORMAT FOR SUBMITTING LIVESTOCK DISEASE OUTBREAK DATA TO NIVEDI. (REVISED REPORT-11/07/2018) | | | | | | | | | | | | |
|--|---------------------------------------|--|----------------------|-------------------------------|-------------------|------|-------|---------------------|--------------------|-----------------|------------------|-----------------------|
| NAME OF THE COLLABORATING UNIT | | ADDRESS OF THE COLLABORATING UNIT | | | | | | | | | | |
| PI-ACRP-ADMAS of NIVEDI SHIMLA, HIMACHAL PRADESH | | PI-ACRP-ADMAS of NIVEDI-cum-Deputy Director Epidemiology, State Veterinary Hospital Complex Cart Road Shimla-171001, Phone: 0177-2650938, 94180-61610 Email: munish_betta@hotmail.com; greckchitender@yahoo.com | | | | | | | | | | |
| REPORT FOR THE MONTH OF | | DATE OF REPORT | | | | | | | | | | |
| July 2019 | | 20.08.2019 | | | | | | | | | | |
| Name of the village* | Latitude and longitude of the village | Postal pin code of the village | Name of the district | Name of the disease affected* | Species affected* | Year | Month | Number of outbreaks | Number susceptible | Number attacked | Number of deaths | Number of vaccination |
| Zunig | 30.6458° N 77.4788° E | 172022 | Shimla | Sheep Pox | Sheep | 2019 | July | 1 | 300 | 43 | 0 | 259 |
| Chaura | 31.5581429° N 77.9467311° E | 172101 | Kinnar | Sheep Pox | Sheep | 2019 | July | 1 | 200 | 40 | 5 | 160 |

*If you know the exact place of the outbreak kindly specify the species i.e., cattle, buffalo and buffaloes and similarly write individually for goats and sheep.

Co-PI-ACRP-ADMAS of NIVEDI
State Veterinary Hospital Complex
Shimla-171001

PI-ACRP-ADMAS of NIVEDI

ASSAM REPORT AUGUST-2020

NIVEDI Prediction

District wise Livestock Disease forewarning for August 2020: Assam

| Districts of Assam | Livestock Diseases | | | | | | | | | | | | |
|---------------------|--------------------|------------|-----|----|-----|-------------|-----|-----|-----|---------|-----|--------------|-----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | IIS | PPR | S&G Pox | SF | Theileriosis | Trypanosomiasis |
| Baksa | NR | NR | VHR | NR | NR | NR | VHR | VLR | NR | NR | NR | NR | NR |
| Barpeta | NR | NR | VHR | NR | NR | NR | VLR | VHR | NR | VHR | NR | NR | NR |
| Bongaigaon | NR | NR | VHR | NR | NR | NR | VHR | VHR | NR | VHR | NR | NR | NR |
| Cachar | NR | NR | VHR | NR | NR | VHR | NR | VHR | NR | NR | NR | NR | NR |
| Chirang | NR | NR | VHR | NR | NR | NR | NR | IR | NR | NR | NR | NR | NR |
| Darrang | NR | NR | VHR | NR | NR | NR | NR | IR | VHR | NR | NR | NR | NR |
| Dhemaji | NR | NR | VHR | NR | NR | VHR | NR | VHR | NR | VHR | NR | NR | NR |
| Dhubri | NR | NR | VHR | NR | NR | VHR | VLR | VHR | IR | NR | NR | NR | NR |
| Dibrugarh | NR | NR | VHR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR |
| Dima Hasao | NR | NR | VLR | NR | NR | NR | VLR | VLR | VLR | NR | NR | NR | NR |
| Goalpara | NR | NR | VHR | NR | VHR | NR | VLR | VHR | VLR | NR | VHR | NR | NR |
| Golaghat | NR | NR | VLR | NR | NR | NR | VLR | VHR | NR | VHR | NR | NR | NR |
| Hailakandi | NR | NR | MR | NR | NR | NR | NR | VLR | NR | NR | NR | NR | NR |
| Jorhat | NR | NR | VHR | NR | NR | VHR | NR | VLR | NR | VHR | NR | NR | NR |
| Kamrup | NR | NR | VLR | NR | NR | VLR | VLR | VLR | IR | VHR | NR | NR | NR |
| Kamrup Metropolitan | NR | NR | VHR | NR | NR | VHR | VLR | VHR | IR | NR | VHR | VHR | NR |
| Karbi Anglong | NR | NR | VLR | NR | VHR | NR | VLR | VLR | NR | VHR | NR | NR | NR |
| Karimganj | NR | NR | VHR | NR | NR | NR | VHR | VLR | NR | NR | NR | NR | NR |
| Kokrajhar | NR | NR | VHR | NR | VHR | NR | VLR | VLR | NR | VHR | NR | NR | NR |
| Lakhimpur | NR | NR | VLR | NR | VHR | VHR | NR | VLR | VLR | NR | VHR | NR | NR |
| Morigaon | NR | NR | VHR | NR | NR | NR | NR | VHR | VLR | NR | VHR | NR | NR |

ASSAM Risk Prediction of Theileriosis for the month of August 2020

Legend:

- Very High Risk (Dark Red)
- High Risk (Red)
- Medium Risk (Orange)
- Low Risk (Yellow)
- Very Low Risk (Light Yellow)
- No Risk / No Data (White)

Assam Report August-2020

Monthly Reports_June_July_August 2020_Guahati Centre

1 message

Dr Durbar Prasad Bora <dprbora@gmail.com>

Dear Sir
I am sending herewith the monthly reports for the months of June, July and August, 2020.
Kindly acknowledge the receipt of the same.

Best Regards
Durbar P Bora
PI, AICRP on ADMAS
Guwahati Centre

Dr Durbar Prasad Bora MVSc, PhD (VRI)
Assistant Professor (Sr. Scale)
Department of Microbiology
College of Veterinary Sciences
Assam Agricultural University
Jorhat-781022
ASSAM (INDIA)
Phn +919654533400
+919654533401
Researchgate: https://www.researchgate.net/profile/Durbar_Bora/
ORCID ID: https://orcid.org/0000-0002-5629-7929
Google Scholar ID: 70038989722
Web of Science ResearcherID: AAE-4100-2020

3 attachments

- Monthly report_August2020_ICAR-ADMAS.doc 5177K
- Monthly report_June2020_ICAR-ADMAS.doc 5178K
- Monthly report_July 2020_ICAR-ADMAS.doc 5178K

Project Directorate on (PD_ADMAS)
Animal Disease Monitoring and Surveillance
INDIAN COUNCIL OF AGRICULTURAL RESEARCH
Hobba, Bangalore - 560023

Name of the AICRP on ADMAS centre
Address of the AICRP on ADMAS Centre
Report for the month of
Date of report

Assam Agricultural University, Khanapara, Guwahati Centre
College of Veterinary Science, Khanapara, Guwahati-781022
August, 2020
05.10.2020

| Name of the village | Latitude and longitude of the village | Postal pin code of the village | Name of the district | Name of the disease | Species affected * | Year | Month | Number of outbreaks | Number susceptible | Number attacked | Number of deaths | Number vaccinated |
|---------------------|---------------------------------------|--------------------------------|----------------------|---------------------|--------------------|------|--------|---------------------|--------------------|-----------------|------------------|-------------------|
| VIII-Dalonghat | 26°7'1N 92°9'0 E | 784528 | Darrong | Paramphistomiasis | Cattle | 2020 | August | 1 | 1500 | 8 | Nil | Nil |
| VIII-Christianbasti | 26°15' N 91°7' E | 781005 | Kamrup (M) | Babesiosis | Dog | 2020 | August | 1 | 40 | 2 | 1 | Nil |
| Baruahchariali | 26°75' N 94°20' E | 785001 | Jorhat | Babesiosis | Dog | 2020 | August | 1 | 40 | 2 | 1 | Nil |
| Mirza | 26°29' N 91°69' E | 781125 | Kamrup (M) | Theileriosis | Cattle | 2020 | August | 1 | 700 | 2 | - | Nil |
| Halogaon | 26°14' N 91°73' E | 781103 | Kamrup (M) | canine distemper | Local birds | 2020 | August | 1 | 700 | 20 | 12 | Nil |

(D P Bora)

KARNATAKA REPORT SEPTEMBER-2021



ProMED

INTERNATIONAL SOCIETY
FOR INFECTIOUS DISEASES

Published Date: 2021-09-30 11:40:53

Subject: PRO/SOAS> Foot & mouth disease - India (10): (Karnataka) cattle

Archive Number: 20210930.8698795

FOOT & MOUTH DISEASE - INDIA (10): (KARNATAKA) CATTLE

A ProMED-mail post

<http://www.promedmail.org>

ProMED-mail is a program of the
International Society for Infectious Diseases
<http://www.isid.org>

Date: Fri 24 Sep 2021 18:33 IST

Source: The Hindu [edited]

<https://www.thehindu.com/news/national/karnataka/foot-and-mouth-disease-reported-in-hassan/article36652024.ece>

District wise Livestock Disease Risk Forewarning for September 2021: Karnataka

| Districts of Karnataka | Livestock Diseases | | | | | | | | | | | | |
|------------------------|--------------------|------------|-----|-----|-----|-------------|-----|-----|-----|---------|----|--------------|----------------|
| | Anthrax | Babesiosis | BQ | BT | ET | Fasciolosis | FMD | HS | PPR | S&G Pox | SF | Theileriosis | Trypanosomosis |
| Gulbarga | NR | NR | VLR | VLR | NR | NR | VLR | VLR | NR | VHR | NR | NR | NR |
| Hassan | NR | NR | VHR | VLR | NR | NR | VHR | VLR | VLR | VHR | NR | NR | NR |
| Haveri | VHR | NR | VLR | NR | NR | NR | HR | HR | NR | HR | NR | NR | NR |
| Kodagu | NR | NR | HR | NR | NR | NR | VHR | VLR | NR | NR | NR | NR | NR |
| Kolar | NR | NR | NR | VLR | NR | NR | HR | VLR | NR | NR | NR | NR | NR |
| Koppal | VHR | NR | VLR | VLR | NR | NR | LR | MR | VLR | NR | NR | NR | NR |
| Mandya | NR | NR | NR | VLR | NR | NR | VHR | VLR | NR | NR | NR | NR | NR |
| Mysore | NR | NR | VHR | NR | NR | NR | VHR | VLR | VLR | NR | NR | NR | NR |
| Raichur | VHR | NR | VLR | NR | NR | NR | VLR | LR | VLR | NR | NR | NR | NR |
| Ramanagara | NR | NR | NR | VLR | NR | NR | VHR | NR | VLR | NR | NR | NR | NR |
| Shimoga | NR | NR | VHR | NR | VHR | NR | VHR | MR | VLR | NR | NR | NR | NR |
| Tumkur | VHR | NR | HR | VLR | NR | NR | VHR | VHR | VLR | NR | NR | NR | NR |
| Udupi | NR | NR | NR | NR | NR | NR | HR | NR | NR | NR | NR | NR | NR |
| Uttara Kannada | NR | NR | NR | NR | NR | NR | HR | VLR | NR | NR | NR | NR | NR |
| Yadgir | NR | NR | VLR | VLR | NR | NR | VLR | VLR | VLR | NR | NR | NR | NR |

The outbreak of foot-and-mouth disease (FMD) among cattle in Hassan has left both farmers and veterinarians worried.

Cases have been reported from Arkalgud, Arsikere, Channarayapatna, and Sakleshpur taluks. While the farming community, depending on cattle for regular income, are worried about their livelihood, the veterinarians are struggling hard to provide treatment, amidst a shortage of staff members.

The Department of Veterinary and Animal Husbandry has reported over 150 cases in the district so far. As of [Fri 24 Sep 2021], the animals in 16 villages of the district are being treated. So far the death of one animal had been reported. However, farmers claim more animals have died over the last month. The death of cattle has an impact on milk production, affecting milk producers.

The vaccination for cattle is done once in six months under the National Animal Disease Control Programme. However, the vaccination drive was not done in the last year, owing to the COVID-19 pandemic. "The outbreak of foot and mouth disease is due to the failure of the government in conducting the vaccination. If the vaccination had been done as per the schedule, farmers would not have suffered", H Yoga Ramesh, president of Potato Club in Arkalgud, told media.

Following reports of the disease, veterinarians have been treating affected animals. "Against 24 sanctioned posts of veterinary doctors in Arkalgud taluk, we are only five people. Every doctor is in charge of 2 or more hospitals. We are struggling hard to treat animals", said a veterinarian.

Unless the vacancies were filled up the department cannot deliver services fully. There was a shortage of staff members of other cadres as well, he added.

Considering the cases, the department has launched a ring vaccination programme. Under this, the animals in a 5-kilometre [3.1 mi] radius of the village, where the disease breakout was reported, would be vaccinated. "We have sufficient stock for the ring vaccination. We are planning a mass vaccination in October [2021], where we will cover 6.58 lakh [658 000] animals," said KR Ramesh, Deputy Director of Veterinary and Animal Husbandry in Hassan.

...

Communicated by:

ProMED-SoAs from HealthMap Alerts
<promed-SoAs@promedmail.org>

9.1 Correlational Assessment

The number of outbreaks predicted and outbreaks actually reported were reported in table 9.1. It has been observed from the table that outbreaks predicted two months in advance and timely alerts were issued that helped the stakeholders to take appropriate preventive measures with in time and accordingly the reported outbreaks were very less. Though the use of artificial intelligence system is more beneficial for accurately predicting the livestock disease outbreaks, there are yet number of limitations, namely, there are expected to be under reporting and also non-reporting cases which created the uncertainties in the model predictions while translating model inputs in to model outputs. However, identifying these uncertainties in the prediction using statistical models and highlighting the importance of quality data may improve the model accuracy and confidence while building the model for livestock disease forecasting.

Table 9.1: Number of districts predicted for livestock diseases risk events and reported outbreaks

| Sl No | Livestock diseases | September-2020 | | October-2020 | |
|-------|-----------------------------------|--|--|--|--|
| | | No. of districts predicted for the disease risk events | No. of districts reported the disease outbreaks* | No. of districts predicted for the disease risk events | No. of districts reported the disease outbreaks* |
| 1 | Anthrax | 40 | 8 | 34 | 12 |
| 2 | Babesiosis | 34 | 12 | 36 | 7 |
| 3 | Black quarter | 59 | 18 | 52 | 12 |
| 4 | Bluetongue | 1 | NA | NA | NA |
| 5 | Enterotoxaemia | 16 | 6 | 20 | 4 |
| 6 | Fasciolosis | 48 | 29 | 59 | 14 |
| 7 | Foot and mouth disease | 92 | 13 | 104 | 26 |
| 8 | Haemorrhagic septicaemia | 81 | 22 | 52 | 6 |
| 9 | Peste des petits ruminants | 56 | 16 | 46 | 15 |
| 10 | Sheep & Goat pox | 27 | 2 | 32 | 9 |
| 11 | Swine fever | 34 | 8 | 31 | 4 |
| 12 | Theileriosis | 33 | 17 | 39 | 12 |
| 13 | Trypanosomosis | 39 | 13 | 42 | 17 |

*Which takes in to account of action taken for prediction and non-reporting of cases

10. Launch of Mobile Android app&link to download

Livestock forewarning application (LDF) can be downloaded following the link provided:http://www.nivedi.res.in/android_nadres/LDF.apk and google play store link also provided <https://play.google.com/store/apps/details?id=info.androidhive.ldf>


Further launch of LDF application was done, the news provided below.

The collage consists of several news articles and social media posts from various Indian publications:

- United News of India:** Headline: "Mobile app forewarning of Livestock diseases launched". Description: "The app will alert farmers about diseases that could affect their animals, thus giving them time to take preventive measures and reduce losses."
- INDIAN EXPRESS:** Headline: "Mobile app forewarning of Livestock diseases launched". Description: "The app will alert farmers about diseases that could affect their animals, thus giving them time to take preventive measures and reduce losses."
- krishijagran.com:** Headline: "Livestock Disease Forecasting - Mobile Application (LDF - Mobile App)". Description: "The app will alert farmers about diseases that could affect their animals, thus giving them time to take preventive measures and reduce losses."
- the pioneer:** Headline: "NEW APP TO FOREWARN OF DISEASES IN PARM ANIMALS". Description: "The app will alert farmers about diseases that could affect their animals, thus giving them time to take preventive measures and reduce losses."
- Business Standard:** Headline: "Livestock Disease Forecasting - Mobile Application (LDF-Mobile App) Launched". Description: "The app will alert farmers about diseases that could affect their animals, thus giving them time to take preventive measures and reduce losses."
- Outlook:** Headline: "Mobile app forewarning of Livestock disease launched". Description: "The app will alert farmers about diseases that could affect their animals, thus giving them time to take preventive measures and reduce losses."
- Dairy Times:** Headline: "Shri Radha Mohan Singh launches Livestock Disease Forecasting - Mobile Application (LDF-Mobile App)". Description: "The app will alert farmers about diseases that could affect their animals, thus giving them time to take preventive measures and reduce losses."
- Green Ecosystem:** Headline: "LDF - Mobile App". Description: "The app will alert farmers about diseases that could affect their animals, thus giving them time to take preventive measures and reduce losses."

On the right side of the collage, there is a logo for NIVE DI (National Institute of Veterinary Epidemiology and Preventive Medicine) and a photograph of a modern building complex.

Livestock Disease Forewarning (LDF Mobile App)



To extend the reach of the NADRES forewarning bulletin among the various stakeholders, a Mobile Application named Livestock Disease forewarning app "LDF-Mobile App" was developed. The forewarning methodology adapted in the "mobile app" remains the same as monthly bulletin; it provides user interface to know the predicted forewarning results stored in NADRES MySQL database. A PHP web-based service is developed in Java to extract the results of forewarning two months in advance by keying state name, district name and disease name and display the same in the mobile app. In addition to forewarning, the LDF-Mobile App also provides the details of clinical samples to be collected in case of outbreaks of the listed diseases for laboratory confirmation and immediate preventive measures to be taken up in case of positive prediction/disease confirmation. The LDF mobile app is available at ICAR-NIVEDI website. It is available on Google play store.

11. FARMERS EMPOWERMENT THROUGH IT: DISEASE RISK COMMUNICATION (COLLABORATION WITH FRUITS, NIC, GOVT. OF KARNATAKA)

In addition to NADRES V₂ (The National Animal Disease Referral Expert System), ICAR-NIVEDI collaborated with NIC, Govt. of Karnataka, Karnataka State for sending the SMS alerts directly to farmers who have registered in FRUITS (Farmers Registration and Unified Beneficiary Information System). The information alerts on risk prediction of six livestock diseases were sent through SMS to farmers is presented in Table 11.A. During December 2021, a total of **17,17,253** SMS alerts were sent to farmers.

Table 11.A: Number of famers received the SMS alert through FRUITS application during November 2021

| Disease Name | District Name | No. of farmers receive d SMS | Disease Name | District Name | No. of farmers received SMS |
|--------------------|-----------------|------------------------------|--------------|------------------|-----------------------------|
| Anthrax | Bellary | 2830 | BT | Gadag | 12406 |
| | Haveri | 24001 | | Hassan | 80621 |
| | Kolar | 24287 | | Koppal | 14749 |
| | Koppal | 14749 | | Raichur | 5524 |
| | Tumkur | 89353 | | Tumkur | 89361 |
| BQ | Bangalore | 25418 | FMD | Bangalore Urban | 25418 |
| | Bangalore Rural | 36309 | | Bangalore Rural | 36309 |
| | Chikmagalur | 15426 | | Chamarajanagar | 26169 |
| | Chitradurga | 17876 | | Chikkaballapur a | 37795 |
| | Hassan | 80621 | | Chikmagalur | 15425 |
| | Haveri | 24001 | | Dakshina Kannada | 36889 |
| | Mandya | 109624 | | Gadag | 12397 |
| | Mysore | 110182 | | Hassan | 80621 |
| | Raichur | 5524 | | Kolar | 24287 |
| | Shimoga | 34622 | | Koppal | 14749 |
| | Tumkur | 89356 | | Mandya | 109624 |
| | Uttara Kannada | 11665 | | Mysore | 110182 |
| | Bellary | 2830 | | Ramanagara | 57709 |
| | Chikkaballapura | 37795 | | Tumkur | 89353 |
| | Chitradurga | 14876 | | Udupi | 30126 |
| | Davanagere | 24529 | | Uttara Kannada | 11665 |
| Grand Total | | | | | 1717253 |

12. Appendix

a) R Code

```
#parsmonth_number=8; year_number=2006; current_year=2017;  
  
nadres_func=function (current_year, year_number, month_number)  
{  
  
args= commandArgs(trailingOnly=TRUE)  
  
if (length(args)<3) {  
  
stop("Correct number of arguments must be supplied", call.=FALSE)  
  
}  
  
current_year=args[1]  
  
year_number=args[2]  
  
month_number=args[3]  
  
df_total<-NULL  
  
month_name=data.frame(month=c(1:12),  
  
month_names=c("February","February","October","October","May","October","October","October",  
"October","October","January","January")  
  
)  
  
ss<-fread(file="NADRES.csv",header=T,check.names = F)  
  
col_pars=names(ss)  
  
vars= paste(col_pars[7:ncol(ss)],collapse = "+")  
  
options(verbose = F)  
  
for(disease in c(8,10,11,12,24,31,35,37,48,60,62,65,70,72,79))  
{  
  
# disease=8  
  
rs<-dbSendQuery(mydb,"SELECT  
index_state.state_name,index_state.state_id,index_district.district_id, index_district.district_name,
```



year_list.year, outbreak_data_final.month, ls_sp_index.species_name,disease_master.disease_id, disease_master.disease_name, outbreak_data_final.number_of_outbreaks, outbreak_data_final.number_susceptible, outbreak_data_final.number_of_attacks, outbreak_data_final.number_of_deaths

```
FROM ls_sp_index INNER JOIN (year_list INNER JOIN (disease_master INNER JOIN
(index_district INNER JOIN (index_state INNER JOIN outbreak_data_final ON
index_state.state_id= outbreak_data_final.state_id) ON index_district.district_id =
outbreak_data_final.district_id) ON disease_master.disease_id= outbreak_data_final.disease_id)
ON year_list.year = outbreak_data_final.year) ON ls_sp_index.species_id=
outbreak_data_final.species_id; ")
```

```
data = fetch(rs, n=-1)
```

```
# year change
```

```
data<-subset(data,data$year>=year_number&&data$disease_id==disease)
```

```
df<-sqldf("SELECT
state_id,state_name,district_id,district_name,disease_id,disease_name,month,sum(number_of_outbr
eaks)as outbreak FROM data GROUP BY
state_id,district_id,state_name,district_name,month,disease_id,disease_name",drv="SQLite")
```

```
ss1<-subset(ss,ss$disease_id==disease)
```

```
attach(ss1,warn.conflicts = F)
```

```
attach(df,warn.conflicts = F)
```

```
dd<-merge(ss1, df, by = c("state_id","district_id","disease_id","month")),all.x=TRUE)
```

```
attach(dd,warn.conflicts = F)
```

```
out<-data.frame(outbreak)
```

```
out<-ifelse(outbreak>=1,1,0)
```

```
out[is.na(out)]<-0
```

```
final<-cbind(dd,out)
```

```
final1<-final[which(final$disease_id==disease),]
```

```
cat("For disease: ",as.character(unique(ss1[,"disease_name"])), "\n")
```



```

ncs= ncol(final1)-5

temp = data.frame(final1[,8:ncs])

for(i in 1:ncol(temp)){
  temp[is.na(temp[,i]), i] <- mean(temp[,i], na.rm = TRUE)
}

final2<-
  cbind(final1$state_id,final1$state_name.x,final1$district_id,final1$district_name.x,final1$disease_id,final1$disease_name.x,final1$out,final1$month,temp)

setnames(final2,old=c("final1$state_id","final1$state_name.x","final1$district_id","final1$district_name.x","final1$disease_id","final1$disease_name.x","final1$out","final1$month"),new=c("state_id","state_name","district_id","district_name","disease_id","disease_name","out","month"))

formula=paste("out ~",vars)

as.formula(formula)

model<-glm(formula,data= final2, family = binomial(link="logit"),maxit=20)

new<-data.frame(final2[,8:ncol(final2)])

prediction<-predict(model,type="response")

n2=randomForest(as.formula(formula),final2)
prediction_rf<-predict(n2,type="response")

gbm_model=gbm.step(data=final2, gbm.x = 8:ncol(final2), gbm.y = 7, family = "bernoulli",
tree.complexity = 1, learning.rate = 0.01,
  bag.fraction= 0.5, n.trees = 5,keep.fold.fit=T,tolerance.method="fixed" , step.size= 5,n.folds = 10)

prediction_gbm<-predict(gbm_model,n.trees=gbm_model$gbm.call$best.trees,type="response")
prediction=numeric()
for (i in 1:length(prediction_gbm)) {
  # if(prediction_glm[i]>prediction_rf[i])
  # {
  #   if(prediction_glm[i]>prediction_gbm[i])
  #   {
  #     prediction[i]=prediction_glm[i]
  #   }
  #   if(prediction_glm[i] >= prediction_gbm[i] && prediction_glm[i] >= prediction_rf[i])
  #
}

```

```

    prediction[i]=prediction_glm[i];
}

if(prediction_gbm[i] >= prediction_glm[i] && prediction_gbm[i] >= prediction_rf[i])
{
    prediction[i]=prediction_gbm[i];
}

if(prediction_rf[i] >= prediction_glm[i] && prediction_rf[i] >= prediction_gbm[i]) {
    prediction[i]=prediction_rf[i];
}

}

summary(prediction)
vv<-round(prediction,2)

df1<-cbind(final2,vv)

df_total<-rbind(df_total,df1)

gc()

}

f=function(m){
if(m<=0.0) i=1
else if(m>=0.0 && m<=0.20) i=2
else if(m>=0.21 && m<=0.40) i=3
else if(m>=0.41 && m<=0.60) i=4
else if(m>=0.61 && m<=0.80) i=5
else i=6
}

df_total$cate=factor(mapply(f,df_total$vv),levels=1:6,labels=c("", "", "", "MR", "", "HR"))

write.csv(df_total,"nadres_outbreak.csv")

##### ACCURACY

df_total=read.csv("nadres_outbreak.csv",header = T)

```

```

dir.create(path = paste(month_name[month_number,2],current_year))

df_poa=df_total

df_poa$cate=factor(mapply(f,df_poa$vv),levels=1:6,labels=c(0,0,0,0,1,1))

df_poa=df_poa[which(df_poa$month==month_name[month_number,1]),]

df_p=df_poa[,c("disease_name","out","cate")]

df_acc=cbind(data.frame(c(1:ow(df_tot_res))),data.frame(df_tp_tn[,1]),(df_tp_tn[,2]/df_tot_res[,2])*100)

df_acc=setNames(df_acc,c("No","Disease","Accuracy"))

print(df_acc)

dis_acc=paste(paste(month_name[month_number,2]," ",current_year,"/",sep=""),"Disease Accuracy ",month_name[month_number,2]," ",current_year, ".csv",sep="")

write.csv(df_acc,dis_acc,row.names = F)

#####PLOT

i=1

plot_dir=paste(paste(month_name[month_number,2]," ",current_year,"/",sep=""),month_name[month_number,2]," ",current_year," N",sep="")

dir.create(path = plot_dir)

disease = c(8,10,11,12,31,35,37,48,60,65,70,72,79)

while(i<=length(disease))

{

kar=readOGR(dsn = "1shp/2011_Dist.shp",verbose = FALSE)

cols=as.character(unique(df_total[df_total$disease_id==disease[i],"disease_name"]))

df_disease=df_total[which(df_total$month==month_name[month_number,1]&df_total$disease_id==disease[i]),]

df_disease=df_disease[,c(2:5,(ncol(df_disease)-1))]
```



```

df_disease=setNames(df_disease,c("ST_CEN_CD","state_name","DT_CEN_CD","district_name","vv"))

kar@data=merge(data.frame(kar@data),data.frame(df_disease),by=c("ST_CEN_CD","DT_CEN_CD"),all.x=T)

kar$vv[is.na(kar$vv)]<-0

#View(kar@data)

colours<-c("#FFFFFF", "#FFFF00", "#FFC1C1", "#FF7150", "#FF8500", "#FF0000")

kar$lb=factor(mapply(f,kar$vv),levels=1:6,labels=c("No Risk / No Data","Very Low Risk","LowRisk","MediumRisk","HighRisk","Very High Risk"))

ICAR
cols=gsub("& ", "and",cols)

disname= gsub("\.", " ",cols)

cat("Plot for disease:",disname,"\\n")

plot_loc=paste(plot_dir,"/",disname,"/",sep="")

dir.create(plot_loc)

file_name=paste(plot_loc,disname,".png",sep="")

plot_title= paste(disname," risk prediction(",month_name[month_number,2],"",
"current_year,)",sep="")

png(file_name)

print(spplot(obj=kar,c("lb"),col.regions=colours,main= plot_title,scales=list(draw = TRUE)))

dev.off()

i=i+1

```



b) Abbreviations

NADRES : National Animal Disease Referral Expert System

R : R environment for statistical computing

BQ : Black Quarter

BT : Bluetongue

ET : Enterotoxaemia

FMD : Foot and Mouth disease

HS : Haemorrhagic Septicaemia

PPR : Peste des Petits Ruminants

SGP : Sheep and Goat pox

SF : Swine Fever

hPa : Hectopascals

NR : No risk/No data available

VLR : Very low risk

LR : Low risk

MR : Moderate risk

HR : High risk

VHR : Very high risk



13. INFECTION WITH SARS-COV-2 IN ANIMALS

Aetiology, Epidemiology, Diagnosis, Prevention and Control

Last updated on 3 July 2020

SARS-CoV-2 is the pathogenic agent that causes the disease COVID-19 and was first reported in December 2019. SARS-CoV-2 is thought to have emerged from an animal source and then spilled-over to the human population. Although genetically closely related viruses have been isolated from *Rhinolophus* bats, the exact source of SARS-CoV-2 and route of introduction into the human population has not been established.

The current pandemic of COVID-19 is being sustained through human-to-human transmission. Animal infections with SARS-CoV-2 have been reported by several countries. Several animal species have proven to be susceptible (Table 1) to infection with SARS-CoV-2 either naturally or by experimental infection. Important livestock species (pigs and poultry) have been demonstrated not to be susceptible to infection through experimental studies. Further studies are needed to understand if and how different animals could be affected by SARS-CoV-2.

It is important to monitor infections in animals to better understand their epidemiological significance for animal health, biodiversity, and human health. Evidence from risk assessments, epidemiological investigations, and experimental studies do not suggest that live animals or animal products play a role in SARS-CoV-2 infection of humans.

Infection with SARS-CoV-2 is not included in the OIE List of Diseases. However, consistent with the reporting obligations of Members outlined in Articles 1.1.4. and 1.1.6. of the OIE *Terrestrial Animal Health Code* relating to emerging diseases, the disease should be notified to the OIE through the OIE's World Animal Health Information System or via email.

AETIOLOGY

Classification of the causative agent

Corona viruses (CoVs) are enveloped, positive-sense, single-stranded RNA viruses. SARS-CoV-2 is a *beta coronavirus*, a genus that includes several coronaviruses (SARS-CoV, MERS-CoV, bat SARS-like CoV, and others) isolated from humans, bats, camels, and other animals.

Susceptibility to physical and chemical action.

SARS-CoV-2 is inactivated by 62–71 % ethanol, 0.5% hydrogen peroxide or 0.1% sodium hypochlorite, within 1 minute, or 0.05–0.2 % benzalkonium chloride or 0.02 % chlorhexidine digluconate.

Survival:

In experimental conditions, SARS-CoV-2 remained viable in the environment after aerosolisation for at least 180 minutes. Experiences with other CoVs such as SARS-CoV, MERS-CoV, or endemic human corona viruses show that:

They can persist on surfaces such as metal, glass or plastic for up to 9 days, but can be efficiently inactivated by surface disinfection procedures, as listed above.

SARS-CoV was found to remain infectious for 14 days at 4 °C, but for only 2 days at 20 °C in sewage water.

Epidemiology

Hosts

Although current evidence suggests that SARS-CoV-2 emerged from an animal source, that source has yet to be identified. The pandemic is driven by person-to-person transmission through respiratory droplets from coughing, sneezing, and talking. Genetic sequence data reveal that SARS-CoV-2 is genetically closely related to other corona viruses circulating in *Rhinolophus* bat (horseshoe bat) populations. To date, there is not enough scientific evidence to identify the source of SARS-CoV-2 or to explain the original route of transmission to humans (which may involve an intermediate host).

Several animal species have tested positive for SARS-CoV-2, mostly as a result of close contact with humans infected with SARS-CoV-2. In addition, preliminary findings from experimental infection studies suggest that poultry and pigs are not susceptible to SARS-CoV-2 infection. The list of animal species for which information on natural or experimental infection is available is presented in Table 13.1.



Table 13.1. Summary of findings in animalstodate

| Species | Type of infection | Susceptibility (none/low/high) | Clinical signs | Transmission |
|--|--------------------------|--|---------------------------------------|--|
| Pigs | Experimental | None | No | No |
| Poultry (chicken, ducks, and turkeys) | Experimental | None | No | No |
| Dogs | Natural and experimental | Low | No (possible in some cases) | No |
| Cats(domestic) | Natural and experimental | High | Yes (none to very mild in some cases) | Yes, between cats |
| Tigers and lions | Natural | High | Yes | Yes, between animals |
| Ferrets | Experimental | High | No (very mild in some cases) | Yes, between ferrets |
| Minks (American minks, <i>Neovison vison</i>) | Natural | High | Yes | Yes, between minks and suggested from mink to humans |
| Egyptian fruit bats (<i>Rousettus aegyptiacus</i>) | Experimental | High | No | Yes, between Fruitbats |
| Golden Syrian hamsters | Experimental | High | Yes (none to very mild in some cases) | Yes, between hamsters |
| Macques (<i>Macaca fascicularis</i> and <i>Macaca mulatta</i>) | Experimental | High | Yes | Yes |

Transmission

Information on the routes of transmission of SARS-CoV-2 among animals is limited. However, as for other respiratory viruses, it appears to be transmitted to animals and between animals by direct contact (e.g droplets). SARS-CoV-2 has been found in secretions from the respiratory tract and in faeces.

Viraemia, incubation and infectious period

In laboratory settings, the incubation period in animals appears to be similar to the one seen in humans (i.e., between 2 and 14 days, with a mean duration of 5 days). However, more studies are required to solidly estimate the mean duration of incubation and the infectious periods.

Sources of virus

The main source of the virus is respiratory droplets and respiratory secretions, although it is possible to isolate SARS-CoV-2 from faeces of infected animals.



Pathogenesis

In laboratory settings, infected animals showed presence of the virus in the respiratory tract and, in some cases, lesions in the trachea and lungs, associated with dyspnoea and cough.

Occurrence and impact

There have been sporadic reports of companion animals and captive wild animals infected with SARS-CoV-2. With regard to production animals, to date, SARS-CoV-2 has only affected mink farms in the Netherlands, with high morbidity and low mortality.

Diagnosis

Knowledge about the susceptibility of different animal species to SARS-CoV-2 infection and clinical signs is, to date, limited (see Table 1).

Clinical diagnosis

Knowledge about clinical disease manifestations in animals is limited. Current evidence suggests clinical signs may include, but are not limited to, coughing, sneezing, respiratory distress, nasal discharge, ocular discharge, vomiting or diarrhoea, fever and lethargy. As in humans, asymptomatic infections occur.

Lesions

More studies are needed to systematically categorise the lesions resulting from infection with SARS-CoV-2 in animals.

In transgenic mice expressing the human version of the SARS-CoV-2 ACE2 receptor, the typical histopathology outcome was interstitial pneumonia with significant inflammatory cell infiltration around the bronchioles and blood vessels, and viral antigens were detected in bronchial epithelial cells and alveolar epithelial cells. These pathological findings were not seen in wild type mice infected with SARS-CoV-2. In golden Syrian hamsters, histopathological changes were reported in the respiratory tract and spleen. Rhesus macaques infected with SARS-CoV-2 presented lesions similar to those seen in humans. Juvenile cats infected with SARS-CoV-2 presented massive lesions in the

nasal and tracheal mucosa epithelia, and lungs. SARS-CoV-2 can replicate in the upper respiratory tract of ferrets without causing severe disease and only resulting in pathological findings such as severe lymphoplasmacytic perivasculitis and vasculitis, increased numbers of type II pneumocytes, macrophages, and neutrophils in the alveolar septa and alveolar lumen, and mild peri bronchitis in the lungs.

Differential diagnosis

All other causes for respiratory or digestive illness should be excluded before a tentative diagnosis for infection with SARS-CoV-2 is made. Existence of an epidemiological link with a confirmed infection in humans or other animals should be considered when narrowing down the list of differential diagnoses.

Laboratory confirmatory tests are necessary for a final diagnosis.



Laboratory diagnosis

Samples

Depending on the type of test, samples may include single or combinations of oropharyngeal, nasal, and rectal swabs, and blood. Faecal samples may be used in situations where direct sampling is not possible due to risks to the animal or testing staff. Tests should be validated for the purpose, species and matrix to be analysed.

Procedures

Agent identification

- Reverse-transcription polymerase chain reaction (RT-PCR);
- Reverse transcription loop-mediated isothermal amplification (RT-LAMP);
- Other molecular tests developed for use in humans;
- Virus isolation;
- Virus genome sequencing.

Detection of immune response:

- ELISA antibody test;
- Virus neutralisation test (VNT);
- Several other tests for antibody detection.

Prevention and control

Biosecurity and hygiene measures are key to preventing transmission of SARS-CoV-2.

People who are suspected or confirmed to be infected with SARS-CoV-2 should restrict contact with mammalian animals, including pets, just like they would with people during their illness.

Animals suspected or confirmed to be infected with SARS-CoV-2 should remain separated from other animals and humans.

References

1. World Health Organization, (consulted on 11/05/2020)
<https://www.who.int/emergencies/diseases/novel-coronavirus-2019>
2. M. Denis, V. Vanderweerd, R. Verbeeke, A. Laudisoit, L. Wynants, D. Van Der Vliet (2020). COVIPENDIUM: information available to support the development of medical counter measures and interventions against COVID-19 (Version 2020-05-05). Transdisciplinary Insights. <http://doi.org/10.5281/zenodo.3782325>
3. QuestionsandAnswersonCOVID19,(consultedon11/05/2020),
<https://www.oie.int/en/scientific-expertise/specific-information-and-recommendations/questions-and-answers-on-2019novel-coronavirus/>
4. Considerations for sampling, testing, and reporting of SARS-CoV-2 in animals, (consulted on 11/05/2020),
https://www.oie.int/fileadmin/Home/eng/Our_scientific_expertise/docs/pdf/COV-19/Sampling_Testing_and_Reportin_g_of_SARS-CoV-2_in_animals_final_7May_2020.pdf
5. Cohen J. (2020). From mice to monkeys, animals studied for corona virus answers. *Science*, Vol.368, Issue 6488; pp.221-222 <https://science.sciencemag.org/content/368/6488/221>
6. CDC, Corona virus Disease 2019 (COVID-19)–pets & other animals (consulted on 29/05/2020) <https://www.cdc.gov/coronavirus/2019-ncov/daily-life-coping/positive-pet.html>



EPIDEMIOLOGY OF COVID-19

| Epidemiology of COVID-19 ALL INDIA | | | | | | | | | | | |
|---|---|--------------------------------|-----------------------------|------|-------------------|------------------------------------|---|----------------------------------|----------------------------|------------------------------|----------------------------|
| Number of Infections (5 laks Increment) | No of days taken to reach since 22 Jan-2020 | date reached since 22 Jan-2020 | Cumulative Number of deaths | CFR | avg. daily deaths | R ₀ for confirmed cases | Required herd Immunity (Threshold) R ₀ | Total Vaccine Administered (cum) | % of Immunity by Infection | % of immunity by vaccination | total % of Immunity gained |
| 1st 5 Lakh Cases | 156 days | 26-06-20 | 15685 | 3.06 | 116.0 | 1.872 | 46.58 | | 0.04 | | 0.04 |
| Cum 10 Lakh Cases | 20 days | 16-07-20 | 25602 | 2.49 | 481.1 | 1.802 | 44.51 | | 0.07 | | 0.07 |
| Cum 15 Lakh Cases | 12 days | 28-07-20 | 34193 | 2.23 | 711.5 | 1.762 | 43.25 | | 0.11 | | 0.11 |
| cum 20 Lakh Cases | 9 days | 06-08-20 | 41585 | 2.04 | 809.4 | 1.732 | 42.26 | | 0.14 | | 0.14 |
| cum 25 Lakh Cases | 8 days | 14-08-20 | 49036 | 1.97 | 932.2 | 1.732 | 42.26 | | 0.18 | | 0.18 |
| cum 30 Lakh Cases | 8 days | 22-08-20 | 56706 | 1.90 | 964.3 | 1.722 | 41.93 | | 0.22 | | 0.22 |
| cum 35 Lakh Cases | 7 days | 29-08-20 | 63498 | 1.82 | 973.0 | 1.702 | 41.25 | | 0.25 | | 0.25 |
| cum 40 Lakh Cases | 6 days | 04-09-20 | 69561 | 1.74 | 996.3 | 1.692 | 40.9 | | 0.29 | | 0.29 |
| cum 45 Lakh Cases | 6 days | 10-09-20 | 76271 | 1.70 | 1111.0 | 1.692 | 40.9 | | 0.32 | | 0.32 |
| cum 50 Lakh Cases | 5 days | 15-09-20 | 82066 | 1.64 | 1157.6 | 1.692 | 40.9 | | 0.36 | | 0.36 |
| cum 55 Lakh Cases | 6 days | 21-09-20 | 88935 | 1.62 | 1146.0 | 1.692 | 40.9 | | 0.39 | | 0.39 |
| cum 60 Lakh Cases | 6 days | 27-09-20 | 95542 | 1.59 | 1101.5 | 1.692 | 40.9 | | 0.43 | | 0.43 |
| cum 65 Lakh Cases | 6 days | 03-10-20 | 101782 | 1.57 | 1039.3 | 1.692 | 40.9 | | 0.47 | | 0.47 |

| | | | | | | | | | | | |
|--------------------|---------|----------|--------|------|--------|-------|-------|-----------|------|------|-------|
| cum 70 Lakh Cases | 7 days | 10-10-20 | 108334 | 1.55 | 937.0 | 1.702 | 41.25 | | 0.50 | | 0.50 |
| cum 75 Lakh Cases | 8 days | 18-10-20 | 114610 | 1.53 | 784.5 | 1.722 | 41.93 | | 0.54 | | 0.54 |
| cum 80 Lakh Cases | 10 days | 28-10-20 | 120527 | 1.51 | 591.7 | 1.752 | 42.92 | | 0.57 | | 0.57 |
| cum 85 Lakh Cases | 10 days | 07-11-20 | 126121 | 1.48 | 559.3 | 1.772 | 43.57 | | 0.61 | | 0.61 |
| cum 90 Lakh Cases | 12 days | 19-11-20 | 132162 | 1.47 | 503.2 | 1.972 | 49.29 | | 0.65 | | 0.65 |
| cum 95 Lakh Cases | 13 days | 02-12-20 | 138648 | 1.46 | 498.9 | 2.332 | 57.12 | | 0.68 | | 0.68 |
| cum 100 Lakh Cases | 16 days | 18-12-20 | 145136 | 1.45 | 405.3 | 3.083 | 67.56 | | 0.72 | | 0.72 |
| cum 105 Lakh Cases | 26 days | 13-01-21 | 151727 | 1.45 | 253.6 | 2.673 | 62.59 | | 0.75 | | 0.75 |
| cum 110 Lakh Cases | 39 days | 21-02-21 | 156385 | 1.42 | 119.4 | 2.052 | 51.27 | 10651012 | 0.79 | 0.76 | 1.55 |
| cum 115 Lakh Cases | 25 days | 18-03-21 | 159370 | 1.39 | 119.4 | 2.092 | 52.2 | 35923500 | 0.83 | 2.58 | 3.40 |
| cum 120 Lakh Cases | 10 days | 28-03-21 | 161843 | 1.35 | 247.5 | 2.202 | 54.59 | 55180875 | 0.86 | 3.96 | 4.82 |
| cum 125 Lakh Cases | 7 days | 04-04-21 | 165101 | 1.32 | 464.4 | 2.202 | 54.59 | 76405697 | 0.90 | 5.48 | 6.38 |
| cum 130 Lakh Cases | 4 days | 08-04-21 | 167642 | 1.29 | 640.5 | 2.112 | 52.65 | 91881530 | 0.93 | 6.59 | 7.53 |
| cum 135 Lakh Cases | 3 days | 11-04-21 | 170179 | 1.26 | 838.0 | 2.012 | 50.3 | 102000401 | 0.97 | 7.32 | 8.29 |
| cum 140 Lakh Cases | 3 days | 14-04-21 | 173123 | 1.24 | 981.0 | 1.902 | 47.42 | 111913288 | 1.00 | 8.03 | 9.04 |
| cum 145 Lakh Cases | 2 days | 16-04-21 | 175649 | 1.21 | 1260.0 | 1.822 | 45.12 | 117305344 | 1.04 | 8.42 | 9.46 |
| cum 150 Lakh Cases | 2 days | 18-04-21 | 178769 | 1.19 | 1560.0 | 1.742 | 42.59 | 121207098 | 1.08 | 8.70 | 9.78 |
| cum 155 Lakh Cases | 2 days | 20-04-21 | 182533 | 1.18 | 1882.0 | 1.701 | 41.21 | 127428887 | 1.11 | 9.15 | 10.26 |
| cum 160 Lakh Cases | 2 days | 22-04-21 | 186920 | 1.17 | 2193.0 | 1.641 | 39.06 | 132754608 | 1.15 | 9.53 | 10.68 |
| cum 165 Lakh Cases | 1 days | 23-04-21 | 189544 | 1.18 | 2624.0 | 1.611 | 37.93 | 135658324 | 1.18 | 9.74 | 10.92 |

| | | | | | | | | | | | |
|--------------------|--------|----------|--------|------|--------|-------|-------|-----------|------|-------|-------|
| cum 170 Lakh Cases | 2 days | 25-04-21 | 195123 | 1.15 | 2789.0 | 1.581 | 36.75 | 139185173 | 1.22 | 9.99 | 11.21 |
| cum 175 Lakh Cases | 1 days | 26-04-21 | 197894 | 1.13 | 2771.0 | 1.561 | 35.94 | 142524947 | 1.26 | 10.23 | 11.48 |
| cum 180 Lakh Cases | 2 days | 28-04-21 | 204832 | 1.14 | 3469.0 | 1.531 | 34.68 | 147053392 | 1.29 | 10.55 | 11.85 |
| cum 185 Lakh Cases | 1 day | 29-04-21 | 208330 | 1.13 | 3498.0 | 1.521 | 34.25 | 149268772 | 1.33 | 10.71 | 12.04 |
| cum 190 Lakh Cases | 1 day | 30-04-21 | 211853 | 1.12 | 3523.0 | 1.511 | 33.82 | 151998107 | 1.36 | 10.91 | 12.27 |
| cum 195 Lakh Cases | 1 day | 01-05-21 | 215542 | 1.11 | 3689.0 | 1.510 | 33.77 | 153626325 | 1.40 | 11.03 | 12.43 |
| cum 200 Lakh Cases | 2 days | 03-05-21 | 222408 | 1.11 | 3433.0 | 1.501 | 33.38 | 156082136 | 1.44 | 11.20 | 12.64 |
| cum 205 Lakh Cases | 1 days | 04-05-21 | 226188 | 1.10 | 3780.0 | 1.491 | 32.93 | 157750752 | 1.47 | 11.32 | 12.79 |
| cum 210 Lakh Cases | 1 days | 05-05-21 | 230168 | 1.10 | 3980.0 | 1.491 | 32.93 | 159931238 | 1.51 | 11.48 | 12.98 |
| cum 215 Lakh Cases | 2 days | 07-05-21 | 238270 | 1.10 | 4051.0 | 1.481 | 32.48 | 165190000 | 1.54 | 11.86 | 13.40 |
| cum 220 Lakh Cases | 1 day | 08-05-21 | 242347 | 1.10 | 4077.0 | 1.481 | 32.48 | 167493857 | 1.58 | 12.02 | 13.60 |
| cum 225 Lakh Cases | 1 day | 09-05-21 | 246116 | 1.09 | 3769.0 | 1.471 | 32.02 | 168304868 | 1.61 | 12.08 | 13.69 |
| cum 230 Lakh Cases | 2 days | 11-05-21 | 254197 | 1.11 | 4040.0 | 1.471 | 32.04 | 173862643 | 1.65 | 12.48 | 14.13 |
| cum 235 Lakh Cases | 1 day | 12-05-21 | 258317 | 1.10 | 4120.0 | 1.461 | 31.58 | 176045577 | 1.69 | 12.63 | 14.32 |
| cum 240 Lakh Cases | 1 day | 13-05-21 | 262317 | 1.09 | 4000.0 | 1.461 | 31.58 | 178361846 | 1.72 | 12.80 | 14.52 |
| cum 245 Lakh Cases | 2 days | 15-05-21 | 270284 | 1.10 | 3983.0 | 1.461 | 31.58 | 181544536 | 1.76 | 13.03 | 14.79 |
| cum 250 Lakh Cases | 2 days | 17-05-21 | 278719 | 1.11 | 4217.0 | 1.461 | 31.58 | 183817204 | 1.79 | 13.19 | 14.98 |
| cum 255 Lakh Cases | 2 days | 19-05-21 | 287122 | 1.13 | 4201.0 | 1.451 | 31.10 | 186410600 | 1.83 | 13.38 | 15.21 |
| cum 260 Lakh Cases | 2 days | 21-05-21 | 295525 | 1.14 | 4202.0 | 1.451 | 31.10 | 189344105 | 1.87 | 13.59 | 15.45 |
| cum 265 Lakh Cases | 2 days | 23-05-21 | 303720 | 1.15 | 4098.0 | 1.451 | 31.10 | 191877460 | 1.90 | 13.77 | 15.67 |

| | | | | | | | | | | | |
|--------------------|---------|----------|--------|------|--------|-------|-------|-----------|------|-------|-------|
| cum 270 Lakh Cases | 2 days | 25-05-21 | 311388 | 1.15 | 3834.0 | 1.451 | 31.10 | 196463495 | 1.94 | 14.10 | 16.04 |
| cum 275 Lakh Cases | 2 days | 27-05-21 | 318895 | 1.16 | 3754.0 | 1.451 | 31.10 | 201438120 | 1.97 | 14.46 | 16.43 |
| cum 280 Lakh Cases | 3 days | 30-05-21 | 329100 | 1.18 | 3402.0 | 1.451 | 31.10 | 208907723 | 2.01 | 14.99 | 17.00 |
| cum 285 Lakh Cases | 4 days | 03-06-21 | 340702 | 1.20 | 2901.0 | 1.451 | 31.10 | 219831571 | 2.05 | 15.78 | 17.83 |
| cum 290 Lakh Cases | 5 days | 08-06-21 | 344082 | 1.22 | 2565.0 | 1.451 | 31.10 | 223642281 | 2.08 | 16.86 | 18.94 |
| cum 295 Lakh Cases | 5 days | 13-06-21 | 374305 | 1.27 | 4155.0 | 1.451 | 31.10 | 250656362 | 2.12 | 17.99 | 20.10 |
| cum 300 Lakh Cases | 9 days | 22-06-21 | 390660 | 1.30 | 1817.0 | 1.461 | 31.58 | 289961746 | 2.15 | 20.81 | 22.96 |
| cum 305 Lakh Cases | 10 days | 02-07-21 | 401050 | 1.31 | 1039.0 | 1.471 | 32.04 | 339447068 | 2.19 | 24.36 | 26.55 |
| cum 310 Lakh Cases | 13 days | 15-07-21 | 412531 | 1.33 | 884.0 | 1.481 | 32.50 | 390286930 | 2.22 | 28.01 | 30.23 |
| cum 315 Lakh Cases | 13 days | 28-07-21 | 422662 | 1.34 | 780.0 | 1.481 | 32.50 | 445696580 | 2.26 | 31.99 | 34.25 |
| cum 320 Lakh Cases | 13 days | 10-08-21 | 429179 | 1.34 | 501.0 | 1.502 | 33.40 | 517799432 | 2.30 | 37.16 | 39.46 |
| cum 325 Lakh Cases | 14 days | 24-08-21 | 434756 | 1.34 | 398.0 | 1.511 | 33.82 | 594552135 | 2.33 | 42.67 | 45.00 |
| cum 330 Lakh Cases | 12 days | 05-09-21 | 440752 | 1.34 | 499.0 | 1.521 | 34.25 | 686693339 | 2.37 | 49.28 | 51.65 |
| cum 335 Lakh Cases | 15 days | 20-09-21 | 445385 | 1.33 | 309.0 | 1.511 | 33.82 | 817737078 | 2.40 | 58.69 | 61.09 |
| cum 340 Lakh Cases | 22 days | 12-10-21 | 451189 | 1.33 | 264.0 | 1.511 | 33.82 | 963825324 | 2.44 | 69.17 | 71.61 |

DATASOURCE

1.WORLD WIDE DATA SET: <https://data.humdata.org/dataset/novel-coronavirus-2019-ncov-cases>

2.VACCINATION DATA SET: <https://api.covid19india.org/>

CFR: Case Fatality rate , number of deaths for every 100 cases
 Prepared by Spatial Epidemiology Lab , ICAR-NIVEDI, Bengaluru.

Epidemiology of COVID-19 Kerala State R₀Table

| Number of Infections (1 lakhs Increment) | No of days taken to reach since 14 March-2020 | Date reached since 14 March -2020 | Cumulative Number of deaths | CFR | Avg. daily deaths | R ₀ for confirmed cases | Required herd Immunity (Threshold) R ₀ | Total Vaccine Administered (cum) | % of Immunity by Infection | % of immunity by vaccination | Total % of Immunity gained |
|---|---|-----------------------------------|-----------------------------|------|-------------------|------------------------------------|---|----------------------------------|----------------------------|------------------------------|----------------------------|
| 1st 1 lakh Cases | 182 days | 09-11-2020 | 411 | 0.41 | 2.0 | 1.732 | 42.25 | | 0.14 | | 0.14 |
| Cum 2 Lakh Cases | 20 days | 10-01-2020 | 772 | 0.39 | 18.0 | 1.692 | 40.89 | | 0.28 | | 0.28 |
| Cum 3 Lakh Cases | 12 days | 10-13-2020 | 1047 | 0.35 | 23.0 | 1.672 | 40.18 | | 0.43 | | 0.43 |
| Cum 4 Lakh Cases | 14 days | 10-27-2020 | 1377 | 0.34 | 24.0 | 1.642 | 39.09 | | 0.57 | | 0.57 |
| Cum 5 Lakh Cases | 15 days | 11-11-2020 | 1772 | 0.35 | 26.0 | 1.652 | 39.45 | | 0.71 | | 0.71 |
| Cum 6 Lakh Cases | 19 days | 11-30-2020 | 2245 | 0.37 | 25.0 | 1.702 | 41.24 | | 0.85 | | 0.85 |
| Cum 7 Lakh Cases | 19 days | 12-19-2020 | 2787 | 0.40 | 28.0 | 1.802 | 44.5 | | 0.99 | | 0.99 |
| Cum 8 Lakh Cases | 20 days | 01-08-2021 | 3258 | 0.41 | 24.0 | 1.862 | 46.29 | | 1.13 | | 1.13 |
| Cum 9 Lakh Cases | 19 days | 01-27-2021 | 3664 | 0.41 | 21.0 | 2.052 | 51.27 | 58323 | 1.28 | 0.08 | 1.36 |
| Cum 10 Lakh Cases | 18 days | 02-14-2021 | 3986 | 0.40 | 18.0 | 2.983 | 66.48 | 340452 | 1.42 | 0.48 | 1.90 |
| Cum 11 Lakh Cases | 33 days | 03-19-2021 | 4468 | 0.41 | 15.0 | 3.363 | 70.27 | 2298455 | 1.56 | 3.26 | 4.82 |
| Cum 12 Lakh Cases | 31 days | 04-16-2021 | 4878 | 0.41 | 13.0 | 3.433 | 70.87 | 5680114 | 1.70 | 8.06 | 9.76 |
| Cum 13 Lakh Cases | 5 days | 04-21-2021 | 5001 | 0.38 | 25.0 | 2.492 | 59.88 | 6353600 | 1.84 | 9.01 | 10.86 |
| Cum 14 Lakh Cases | 4 days | 04-25-2021 | 5111 | 0.37 | 27.0 | 2.122 | 52.88 | 6847075 | 1.99 | 9.71 | 11.70 |

| | | | | | | | | | | | |
|-------------------|--------|------------|-------|------|-------|-------|-------|----------|------|-------|-------|
| Cum 15 Lakh Cases | 4 days | 04-29-2021 | 5260 | 0.35 | 37.0 | 1.842 | 45.71 | 7229177 | 2.13 | 10.25 | 12.38 |
| Cum 16 Lakh Cases | 2 days | 05-01-2021 | 5357 | 0.33 | 48.0 | 1.762 | 43.24 | 7425416 | 2.27 | 10.53 | 12.80 |
| Cum 17 Lakh Cases | 3 days | 05-04-2021 | 5508 | 0.32 | 50.0 | 1.702 | 41.24 | 7577304 | 2.41 | 10.75 | 13.16 |
| Cum 18 Lakh Cases | 3 days | 05-07-2021 | 5683 | 0.32 | 58.0 | 1.642 | 39.09 | 7869269 | 2.55 | 11.16 | 13.71 |
| Cum 19 Lakh Cases | 2 days | 05-09-2021 | 5815 | 0.31 | 66.0 | 1.612 | 37.95 | 7952556 | 2.69 | 11.28 | 13.97 |
| Cum 20 Lakh Cases | 3 days | 05-12-2021 | 6054 | 0.30 | 80.0 | 1.592 | 37.17 | 8185506 | 2.84 | 11.61 | 14.45 |
| Cum 21 Lakh Cases | 3 days | 05-15-2021 | 6340 | 0.30 | 95.0 | 1.572 | 36.37 | 8417840 | 2.98 | 11.94 | 14.92 |
| Cum 22 Lakh Cases | 3 days | 05-18-2021 | 6613 | 0.30 | 91.0 | 1.562 | 35.96 | 8573567 | 3.12 | 12.16 | 15.28 |
| Cum 23 Lakh Cases | 4 days | 05-22-2021 | 7171 | 0.31 | 139.0 | 1.552 | 35.55 | 8644281 | 3.26 | 12.26 | 15.52 |
| Cum 24 Lakh Cases | 4 days | 05-26-2021 | 7883 | 0.33 | 178.0 | 1.542 | 35.13 | 8789981 | 3.40 | 12.47 | 15.87 |
| Cum 25 Lakh Cases | 4 days | 05-30-2021 | 8642 | 0.35 | 190.0 | 1.542 | 35.13 | 9241456 | 3.55 | 13.11 | 16.65 |
| Cum 26 Lakh Cases | 5 days | 06-04-2021 | 9511 | 0.37 | 174.0 | 1.542 | 35.13 | 10026627 | 3.69 | 14.22 | 17.91 |
| Cum 27 Lakh Cases | 7 days | 06-11-2021 | 10805 | 0.40 | 185.0 | 1.551 | 35.53 | 11102819 | 3.83 | 15.75 | 19.58 |
| Cum 28 Lakh Cases | 9 days | 06-20-2021 | 12061 | 0.43 | 140.0 | 1.561 | 35.94 | 12161549 | 3.97 | 17.25 | 21.22 |
| Cum 29 Lakh Cases | 9 days | 06-29-2021 | 13094 | 0.45 | 115.0 | 1.571 | 36.35 | 13948645 | 4.12 | 19.78 | 23.90 |
| Cum 30 Lakh Cases | 8 days | 07-07-2021 | 14108 | 0.47 | 127.0 | 1.581 | 36.75 | 15120585 | 4.26 | 21.45 | 25.70 |
| Cum 31 Lakh Cases | 7 days | 07-14-2021 | 14938 | 0.48 | 119.0 | 1.601 | 37.54 | 16206261 | 4.40 | 22.99 | 27.38 |
| Cum 32 Lakh Cases | 7 days | 07-21-2021 | 15618 | 0.49 | 97.0 | 1.621 | 38.31 | 17427952 | 4.54 | 24.72 | 29.26 |
| Cum 33 Lakh Cases | 6 days | 07-27-2021 | 16327 | 0.49 | 118.0 | 1.641 | 39.06 | 18942910 | 4.68 | 26.87 | 31.55 |
| Cum 34 Lakh Cases | 5 days | 08-01-2021 | 16838 | 0.50 | 102.0 | 1.641 | 39.06 | 20369496 | 4.82 | 28.89 | 33.71 |

| | | | | | | | | | | | |
|-------------------|--------|------------|-------|------|-------|-------|-------|----------|------|-------|-------|
| Cum 35 Lakh Cases | 5 days | 08-06-2021 | 17516 | 0.50 | 136.0 | 1.661 | 39.80 | 21560396 | 4.96 | 30.58 | 35.55 |
| Cum 36 Lakh Cases | 5 days | 08-11-2021 | 18121 | 0.50 | 121.0 | 1.671 | 40.16 | 22443788 | 5.11 | 31.83 | 36.94 |
| Cum 37 Lakh Cases | 5 days | 08-16-2021 | 18744 | 0.51 | 125.0 | 1.671 | 40.16 | 24620672 | 5.25 | 34.92 | 40.17 |
| Cum 38 Lakh Cases | 5 days | 08-21-2021 | 19429 | 0.51 | 137.0 | 1.671 | 40.16 | 25732732 | 5.39 | 36.50 | 41.89 |
| Cum 39 Lakh Cases | 5 days | 08-26-2021 | 20135 | 0.52 | 141.0 | 1.671 | 40.16 | 27264922 | 5.53 | 38.67 | 44.20 |
| Cum 40 Lakh Cases | 3 days | 08-29-2021 | 20542 | 0.52 | 136.0 | 1.651 | 39.43 | 28152388 | 5.67 | 39.93 | 45.60 |
| Cum 41 Lakh Cases | 4 days | 02-09-2021 | 21150 | 0.52 | 152 | 1.641 | 39.06 | 29423239 | 5.82 | 41.73 | 47.55 |
| Cum 42 Lakh Cases | 3 days | 09-05-2021 | 21497 | 0.52 | 116.0 | 1.641 | 39.06 | 29636101 | 5.96 | 42.04 | 47.99 |
| Cum 43 Lakh Cases | 4 days | 09-09-2021 | 22127 | 0.51 | 157.0 | 1.651 | 39.43 | 30767695 | 6.10 | 43.64 | 49.74 |
| Cum 44 Lakh Cases | 5 days | 14-09-2021 | 22780 | 0.52 | 131.0 | 1.651 | 39.43 | 31874863 | 6.24 | 45.21 | 51.45 |
| Cum 45 Lakh Cases | 5 days | 19-09-2022 | 23592 | 0.52 | 162.0 | 1.651 | 39.43 | 33641007 | 6.38 | 47.72 | 54.10 |
| Cum 46 Lakh Cases | 6 days | 25-09-2023 | 24439 | 0.53 | 141.0 | 1.651 | 39.43 | 35064476 | 6.52 | 49.74 | 56.26 |
| Cum 47 Lakh Cases | | | | | | | | | | | |

CFR: Case Fatality rate , number of deaths for every 100 cases

Prepared by Spatial Epidemiology Lab , ICAR-NIVEDI, Bengaluru.



ICAR - National Institute of Veterinary Epidemiology and Disease Informatics

Customer/Client Feedback Form

Feedback for the Livestock Diseases Risk Forewarning Bulletin of January-2022, Volume 10 and Issue 01

(Please return this duly fill in after receiving the outbreak report of March -2022)

1. Details of the number of districts with diseases reported vs. forecast in your state.

| Sl. No | Disease Name | No. of districts in which outbreaks occurred but not alerted** | Measures taken in case of disease forecasted: Yes or No.** | Any other |
|--------|----------------------------|--|---|-----------|
| 1. | Anthrax | | | |
| 2. | Babesiosis | | | |
| 3. | Black Quarter | | | |
| 4. | Bluetongue | | | |
| 5. | Enterotoxaemia | | | |
| 6. | Fasciolosis | | | |
| 7. | Foot and mouth disease | | | |
| 8. | Haemorrhagic septicaemia | | | |
| 9. | Peste des Petits Ruminants | | | |
| 10. | Sheep & Goat pox | | | |
| 11. | Swine fever | | | |
| 12. | Theileriosis | | | |
| 13. | Trypanosomosis | | | |

**Details may be written here.

2. What are the preventive measures taken in case of predicted outbreaks?



3. How would you rate your satisfaction with the following aspects of the services you have received or accessed?

| Description | Very satisfied | Satisfied | Unsatisfied | Not sure |
|---|--------------------------|--------------------------|--------------------------|--------------------------|
| Quality of services provided | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| Timeliness of alerts received | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| Benefits from forecasting of livestock diseases | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| Your awareness of this service | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

4. Suggestions for further improvement of report.

Sign and Signature with Designation

AICRP centre:

Dated:



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

Agrisearch with a Human touch.



ICAR-National Institute of Veterinary Epidemiology and Disease Informatics (ICAR_NIVEDI),

P. B. No.6450, Yelahanka, Bengaluru-560064

Phone: +91-80-23093111, Fax: +91-80-23093222, E-mail: director.nivedi@icar.gov.in